

**INTERNATIONAL JOURNAL OF FOOD AND
NUTRITIONAL SCIENCES**

IMPACT FACTOR ~ 1.021



Official Journal of IIFANS

SHELF LIFE EXTENSION OF VARIOUS TYPES OF FISH MEAT USING SELECTED MODIFIED ATMOSPHERE PACKAGING (MAP) METHODS, REVIEW

M Abd Elgadir^{1*}, A A Al-Hassan¹, Md. Zaidul Islam Sarker²
and Md. Jahurul Haque Akanda³

*Corresponding Author: M Abd Elgadir, ✉ M.saeed@qu.edu.sa

Received on: 14th August, 2016

Accepted on: 1st June, 2017

Modified Atmosphere Packaging (MAP) is well known and has been used for decades as a popular method for food packaging in with the objectives of prolonging the shelf life. Carbon dioxide, nitrogen and oxygen are the most common gases used in this regard. Carbon dioxide and oxygen are used in MAP because of their great effects as preservative. The shelf life extension using these gases depends on many factors such as appropriate gas composition, the ratio between gas and product volume, storage temperature and hygienic practice of processing and packaging. In this reviews the effect of MAP in inhibiting growth of spoilage as well as pathogenic bacteria in selected fish/fish products. It could be used as a guide for fishery producers and processors when shelf life extension is mentioned.

Keywords: Fish, Shelf life, Modified Atmosphere Packaging (MAP), Microbial growth

INTRODUCTION

Packaging technologies are very important in order to protect products against deteriorative effects. These effects may include physical, biochemical and microbial activities from environmental influences. This involves maintenance of quality in packed food, retardation of spoilage, and extension of shelf-life (Restuccia *et al.*, 2010). Packaging technologies may also involve other functions such as convenience, containment, marketing and communication (Lee, 2010). Modified Atmosphere Packaging (MAP) is considered as an effective food preservation method (Chouliara *et al.*, 2007). MAP is well known method of shelf life extension a variety of foods including fresh meat and poultry (Chouliara *et al.*, 2007). MAP in general can be classified according to Seydim *et al.* (2006) into two main categories: high-oxygen

modified atmospheres and low (including vacuum packaging, CO₂ gas flushing and N₂ gas flushing). Wicklund *et al.* (2006) reported that MAP of red meat for retail sale can prolong the microbiological shelf life when compared with traditional oxygen-permeable overwrap. The acceptance value of vacuum-packaged retail fresh and cooked meat has been low because of its dark reddish purple color (Jayasingh *et al.*, 2001). A combination of vacuum packaging and refrigeration storage is believed to be the most effective method of MAP that currently used for extending shelf life of uncooked meats. There are several research works have been recently reported using many types of meat with different methods of MAP (Yuan *et al.*, 2012; Aida Cachaldora, 2013; Kapetanakou *et al.*, 2014; Meredith, 2014; Sisse Jongberg *et al.*, 2014 and Chenglong Liu, 2014). The

¹ Department of Food Science & Human Nutrition, College of Agriculture & Veterinary Medicine, Qassim University, Al-Qassim, Kingdom of Saudi Arabia.

² Department of Pharmaceutical Technology, Faculty of Pharmacy, International Islamic University Malaysia, Kuantan Campus, 25200 Kuantan, Pahang, Malaysia.

³ School of Industrial Technology, Universiti Sains Malaysia, Minden, 11800 Penang, Malaysia.

objective of this review article is to highlight various methods using MAP on shelf life extension of selected types of meats.

Effect of Biological Changes on Shelf Life

The changes in quality of fish can be controlled by inhibiting microbial growth, enzymatic freshness and biochemical changes during handling and storage. Spoilage in fish depends on chemical components of their muscles as well as their different species. Pereira de Abreu *et al.* (2010) mentioned that those changes together with microbial growth and enzymatic induced activities are involved in muscle degradation. This degradation of muscle structure is thought to be caused by enzymes called proteinases such as cathepsin B, D and M and calpain (Godiksen *et al.*, 2009). Earlier, An *et al.* (1994) reported that cathepsins enzymes possibly caused degradation of myofibrils in Pacific whiting fillets kept at 0 °C. On other research, Pantazi *et al.* (2008) found that from microbial viewpoint, *Pseudomonas* spp. is the main organism responsible for deterioration of Mediterranean swordfish proteins. Continuous hydrolysis of Myosin Heavy Chain (MHC) was observed on of rainbow trout muscle stored in iced for five days which resulted in firmness of the fish muscle (Godiksen *et al.*, 2009). On the other hand, no changes were observed in Mg²⁺-Ca²⁺-ATPase, Ca²⁺-ATPase or Mg²⁺-ATPase of sea bass (*Latescalcalifer*) actomyosin but Mg²⁺-EGTA-ATPase activity gradually increased when the fish stored at refrigeration temperature for 12 days. However, conducted research by Chomnawang *et al.* (2007) revealed that Ca²⁺-ATPase activities of catfish fillet decreased during storage of 15 days at temperature of 4 °C. Benjakul *et al.* (2003) reported that the loss in Ca²⁺-sensitivity of myofibrillar protein could be an indicator of the proteolytic degradation of tropomyosin and modification of actin-myosin interaction by oxidation of sulfhydryl (SH) group of myosin. Thanonkaew *et al.* (2006) found that the color changes of cuttlefish are accompanied with the development of rancid odors during frozen storage. Off-odor in both development in red tilapia (*Oreochromis mossambicus* × *O. niloticus*) and seabass (*Latescalcarifer*) was correlated with lipid oxidation during 15 days of storage at 0 °C. Thiansilakul *et al.* (2010) reported that lipid hydrolysis can occur with the action of enzymes. The majority of lipolysis in most stored fish originates from microorganisms and endogenous enzymes mainly triacyl lipase and phospholipase (Aryee *et al.*, 2007). An increase in pH has been observed with some types of fish partially with

increasing the storage time during iced or chilled storage. Masniyom *et al.* (2002) and Ocano-Higuera *et al.* (2009) found that the pH increased throughout the chilled storage of cazon fish and seabass. The shelf-life of chilled fish is generally limited due to the growth of Gram-negative microorganisms such as *Aeromonas*, *Shewanella putrefaciens* and *Pseudomonas* under aerobic condition (Ravi-Sankar *et al.*, 2008). Proteinases such as and cathepsin D, B and L and calpain play an important role on degradation of fish muscle structure (Godiksen *et al.*, 2009). An *et al.* (1994) reported that cathepsins was the main enzyme caused degradation of myofibrils in Pacific whiting fillets occurred at 0 °C. However, Pantazi *et al.* (2008) found that *Pseudomonas* spp. is the main organism responsible for deterioration of proteins. Masniyom *et al.* (2004) claimed that no changes were observed in Mg²⁺-Ca²⁺-ATPase, Ca²⁺-ATPase or Mg²⁺-ATPase of sea bass (*Latescalcalifer*) actomyosin but gradual increase in Mg²⁺-EGTA-ATPase activity during refrigerated were recorded during storage of 12 days. In another research, Chomnawang *et al.* (2007) reported a decrease in Ca²⁺-ATPase activities of catfish fillet when stored at 4°C for 15 days. These changes in Ca²⁺-sensitivity and ATPase activities could be due to the proteolysis. Microbial spoilage of fish can be caused by the activities of microorganisms. The main microorganisms caused the spoilage on chilled fish are gram-negative bacteria (*Shewanella* spp. and *Pseudomonas* spp.) grown on Gram and Huss (2000). Several spoilage bacteria including *S. putrefaciens* and *Pseudomonas* spp. produce inosine monophosphate or hypoxanthine from inosine. An increase in pH was observed in some fish with increasing the storage time during chilled storage (Masniyom *et al.*, 2002). Gram-positive bacteria such as *Bacillus*, *Micrococcus*, *Clostridium*, *Staphylococcus* spp., *Corynebacterium*, *Streptococcus* and *Brochothrichosphaeta* were found to be the most dominant microorganisms in tropical marine fish (Al Bulushi *et al.*, 2010). Furthermore, microorganism like *Acinetobacter*, *Aeromonas*, *Bacillus*, *Clostridium*, *Escherichia*, *Micrococcus*, *Pseudomonas*, *Salmonella* were described as prolific biogenic amine-forming bacteria (Al Bulushi *et al.*, 2010).

Effects of MAP on Shelf-Life Extension

Modified Atmosphere Packaging (MAP) is currently successfully in commercial application for inhibiting spoilage of different types of fish and their products and extending shelf-life. In many research works (Hansan *et al.*, 2007; Mohan *et al.*, 2008; Hansan *et al.*, 2009; Mohan *et al.*,

2009; Fernández, 2010; Torrieri *et al.*, 2011; Yassoralipour *et al.*, 2012; Macé *et al.*, 2013 and Yesudhason *et al.*, 2014). This could be associated with a number of interrelated factors, such as gas favorable consumer perception of MAP technology and as developments on new food packaging materials (Ashie *et al.*, 1996 and Lee, 2010). An increase in shelf-life of nine days was observed on fresh Mediterranean swordfish kept under vacuum condition compared with aerobic packaging (Pantazi *et al.*, 2008). Nitrogen gas (N₂) was used in MAP as a filler to prevent package collapse because of its low solubility in water and also it inhibited the growth of aerobic spoilage microorganisms and reduced lipid oxidation (Farber, 1991). It was reported that O₂ could be used in low concentrations in fish products to avoid the outbreak of strictly anaerobic pathogens such as nonproteolytic *C. botulinum* (Rutherford *et al.*, 2007). Carbon dioxide (CO₂) commonly becomes more effectively as an antimicrobial agent in fish because of its fungistatic and bacteriostatic properties. It is able to dissolve into the liquid phase in fish muscles and this could be associated with the increased carbonic acid (Banks *et al.*, 1980). The ratio of the volume of gas and volume of food product (G/P ratio) may need to be twice the volume of meat for adequate microbial retardation, although CO₂ is dissolved and absorbed into the meat surface during storage (McMillin, 2008). Ozogual *et al.* (2004) and Goulas *et al.* (2005) stored mussel (*Mytilus galloprovincialis*) under refrigeration temperature using CO₂ in MAP and found an increase in CO₂ in the range of 60-80%, which mainly due to an extension in the lag phase of organisms of the fish. On the other hand, 100% CO₂ enriched atmosphere could lower microbial counts of rainbow trout (Arashisar *et al.*, 2004). CO₂ commonly becomes more effective as an antibacterial agent when its concentration is increased. It retards the microbial growth of spoilage microorganisms such as *Shewanella* spp. and *Pseudomonas* spp. Emborg *et al.* (2005) claimed that microbial growth is generally inhibited by higher carbon dioxide concentration but *P. phosphoreum* was found to be more resistant to CO₂. Moreover, it was found that in higher concentration of CO₂ (80%) the growth of microorganism in mussel was inhibited (Goulas *et al.*, 2005). Lipid oxidation occurs in many types of fish particularly when stored in inappropriate conditions. For instance, during frozen storage color changes of cuttlefish were found to be accompanied with the development of rancid odors (Thanonkaew *et al.*, 2006). Moreover, in both red tilapia (*Oreochromis mossambicus* × *O. niloticus*) and seabass (*Lateolabrax japonicus*) off-odor development was found to be

correlated with lipid oxidation during 15 days of iced storage (Thiansilakul *et al.*, 2010). Aryee *et al.* (2007) reported that majority of lipolysis in most stored fish were as a result of activities of endogenous enzymes and microorganisms. These mainly include triacyl lipase and phospholipase. Earlier, an increase in values of 1, 2- diacylglycerol and Free Fatty Acid (FFA) were observed in minced mackerel during storage at 2-3 °C for 15 days (Hwang and Regenstein, 1993). Young *et al.* (2014) investigated shelf life of both cooked and raw black tiger prawns (*Penaeus monodon*). The samples were cooked-chilled and packed in a combination of gases consisting 40% CO₂, 30% O₂ and 30% N₂ in vacuum pouches for 14 days. It was found that the method of MAP was most efficient at preserving their sensorial and textural properties. At both storage temperatures, this method also increased the lag phase of microbial growth which could be considered as most effective method due to prawns preserved textural, sensorial qualities and lowered microbial counts.

Safety Issues of MAP Used for Fish Packaging

When MAP is used for shelf-life extension of fish, the combination between elevated CO₂ and MAP reduce microbial growth and enzymatic activity. It was found that generally gram negative aerobic bacteria are more resistant in MAP. However, *L. monocytogenes*, *E. coli*, *S. aureus* and *C. botulinum* outbreaks have attracted the attention widely (Kimura *et al.*, 1999). Growth of pathogens such as *Salmonella* spp., and *E. coli* and *L. monocytogenes* were observed widely (Kimura *et al.*, 1999). CO₂ at a level of 80% was found to be significant in reducing growth rate of *L. monocytogenes* in buffered nutrient broth at 7.5 °C (Hendricks and Hotchkiss, 1997). In another study, it was claimed that modified atmosphere using CO₂ (100% CO₂) reduced the growth of *L. monocytogenes* significantly when packs stored at 3 °C compared to vacuum or air packaging (Rutherford *et al.*, 2007). Growth of *E. coli* O157 and *L. monocytogenes* were retarded in the fish packaged under combination of pyrophosphate and MAP but microbial growth did not inhibited completely (Masniyom *et al.*, 2006). Toxin produced by *C. botulinum* considered as a major concern in relation to safety of MAP fish and their products (Ravi-Sankar *et al.*, 2008). Earlier, Cann *et al.* (1984) found salmon and trout inoculated with spore of *C. botulinum* and kept in MAP at 10-20 °C spoiled before they became toxic. Toxin has been found in both vacuum and MAP packed fish fillets prior to the fish being determined as spoiled (Arritt *et al.*, 2007). It was reported that a combination between

storage temperature and MAP played a major role in shelf-life the extension as it was noticed in *C. botulinum* growth, and the toxin production in retail type packages of fresh salmon fillets (Peck *et al.*, 2008).

CONCLUSION

Modified Atmosphere Packaging (MAP) can be used to extend the shelf life of various types of fish and their products. This can be achieved through retarding microbial growth in fish/fish products which leads to a delay in spoilage, decreasing chemical and physical changes and compounds deterioration as well as. However, combined applications of MAP and other treatment are found to be effective methods in quality maintenance and shelf life extension in the presence of good application practice. Therefore, Modified Atmosphere Packaging (MAP) is a very effective food preservation application that can protect fish/fish products from environmental influence.

REFERENCES

- Al Bulushi I M, Poole S E, Barlow R, Deeth H C and Dykes GA (2010), "Speciation of Gram-Positive Bacteria in Fresh and Ambient-Stored Sub-Tropical Marine Fish", *Int. J. Food Microbiol.*, Vol. 138, pp. 32-38.
- Alfaro B and Hernande I (2013), "Evolution of the Indigenous Microbiota in Modified Atmosphere Packaged Atlantic Horse Mackerel (*Trachurus trachurus*) Identified by Conventional and Molecular Methods", *Int. J. Food Microbiol.*, Vol. 67, pp. 117-123.
- An H, Weerasinghe V, Seymour T A and Morrissey M T (1994), "Cathepsin Degradation of Pacific Whiting Surimi Protease", *J. Food Sci.*, Vol. 59, pp. 1013-1017.
- Aryee A N A, Simpson B K and Villalonga R (2007), "Lipase fraction from the Viscera of Grey Mullet (*Mugil cephalus*) Isolation Partial Purification and Some Biochemical Characteristics Enz", *Microbiol. Technol.*, Vol. 40, pp. 394-402.
- Benjakul S, Visessanguan W, Thongkaew C and Tanaka M (2003), "Comparative Study on Physicochemical Changes of Muscle Proteins from Some Tropical Fish During Frozen Storage Food", *Res. Int.*, Vol. 36, pp. 787-795.
- Bono G and Badalucco C (2012), "Combining Ozone and Modified Atmosphere Packaging (MAP) to Maximize Shelf-Life and Quality of Striped Red Mullet (*Mullus surmuletus*) LWT – Food", *Sci. Tech.*, Vol. 47, pp. 500-504.
- Cachaldora A, García G, Lorenzo J M and García-Fontán M C (2013), "Effect of Modified Atmosphere and Vacuum Packaging on Some Quality Characteristics and the Shelf-Life of Morcilla a Typical Cooked Blood Sausage", *Meat Sci.*, Vol. 93, pp. 220-225.
- Chomnawang C, Nantachai K, Yongsawatdigul J, Thawornchinsobut S and Tungkawachara S (2007), "Chemical and Biochemical Changes of Hybrid Catfish Fillet Stored at 4 °C and its Gel Properties", *Food Chem.*, Vol. 103, pp. 420-427.
- Chouliara E, Karatapanis A, Savvaidis I N and Kontominas M G (2007), "Combined Effect of Oregano Essential Oil and Modified Atmosphere Packaging on Shelf-Life Extension of Fresh Chicken Breast Meat, Stored at 40 °C", *Food Microbiol.*, Vol. 24, pp. 607-617.
- Dalgaard P, Gram L and Hans Henrik Huss H H (1993), "Spoilage and Shelf-Life of Cod Fillets Packed in Vacuum or Modified Atmospheres", *Int. J. Food Microbiol.*, Vol. 19, pp. 283-294.
- de Abreu P, Losada P, Maroto J and Cruz J M (2010), "Evaluation of the Effectiveness of a New Active Packaging Film Containing Natural Antioxidants (from Barley Husks) that Retard Lipid Damage in Frozen Atlantic Salmon (*Salmosalar L.*)", *Food Research Int.*, Vol. 43, pp. 1277-1282.
- Emborg J, Laursen B G and Dalgaard P (2005), "Significant Histamine Formation in Tuna (*Thunnus albacares*) at 2 Degrees C-Effect of Vacuum- and Modified Atmosphere-Packaging on Psychrotolerant Bacteria", *Int J. Food Microbiol.*, Vol. 101, pp. 263-279.
- Fernández K, Aspé E and Roeckel M (2010), "Scaling Up Parameters for Shelf-Life Extension of Atlantic Salmon (*Salmosalar*) Fillets Using Superchilling and Modified Atmosphere Packaging", *Food Control.*, Vol. 21, pp. 857-862.
- Godiksen H, Morzel M, Hyldig G and Jessen F (2009), "Contribution of Cathepsins B L and D to Muscle Protein Profiles Correlated with Texture in Rainbow Trout (*Oncorhynchus mykiss*)", *Food Chem.*, Vol. 113, pp. 889-896.

- Goulas AE, Chouliara I, Nessi E, Kontominas M G and Savvaidis IN (2005), "Microbiological Biochemical and Sensory Assessment of Mussels (*Mytilus galloprovincialis*) Stored Under Modified Atmosphere Packaging", *J. Applied Microbiol.*, Vol. 98, pp. 752-760.
- Gram L and Huss H H (2000), "Fresh and Processed Fish and Shellfish in the Microbiological Safety and Quality of Food", B M Lund, A C Baird Parker and G W Gould (Eds.), Chapman and Hall, London, UK.
- Hansan A A, Morkore T, Rudi K, Olsen E and Eie T (2007), "Quality Changes During Refrigerated Storage of MA-Packaged Pre-rigor Fillets of Farmed Atlantic Cod (*Gadus morhua* L.) Using Traditional MAP CO₂ Emitter and Vacuum", *J. Food Sci.*, Vol. 72, pp. 423-430.
- Hansan AA, Morkore T, Rudi K, Rodbotten M, Bjerke F and Eie T (2009), "Quality Changes of Prerigor Filleted Atlantic Salmon (*Salmo salar* L.) Packaged in Modified Atmosphere Using CO₂ Emitter Traditional MAP and Vacuum", *J. Food Sci.*, Vol. 74, pp. 242-249.
- Hwang K T and Regenstein J M (1993), "Characteristics of Mackerel Mince Lipid Hydrolysis", *J. Food Sci.*, Vol. 58, pp. 79-83.
- Jayasingh P, Cornforth D P, Carpenter C E and Whittier D (2001), "Evaluation of Carbon Monoxide Treatment in Modified Atmosphere Packaging or Vacuum Packaging to Increase Color Stability of Fresh Beef", *Meat Sci.*, Vol. 59, pp. 317-324.
- Jongberg S, Wen J, Tørngren MA and Lund M N (2014), "Effect of High-Oxygen Atmosphere Packaging on Oxidative Stability and Sensory Quality of Two Chicken Muscles During Chill Storage", *Food Packaging and Shelf Life*, Vol. 1, pp. 38-48.
- Kapetanakou A E, Agathaggelou E I and Skandamis P N (2014), "Storage of Pork Meat Under Modified Atmospheres Containing Vapors from Commercial Alcoholic Beverages", *International Journal of Food Microbiol.*, Vol. 178, pp. 65-75.
- Lauzon H L, Stefánsson G, Jónsson G P and Sveinsdóttir K (2002), "Sensory Quality and Shelf Life of MAP Redfish (*Sebastes marinus*) Fillets as Affected by Previous MA Bulk Storage of Whole Fish", *J. Aquatic Food Prod. Tech.*, Vol. 11, pp. 251-273.
- Lee K T (2010), "Quality and Safety Aspects of Meat Products as Affected by Various Physical Manipulations of Packaging Materials", *Meat Sci.*, Vol. 86, pp. 138-150.
- Liu C, Zhang Y, Yang X, Liang R, Mao Y, Hou X, Lu X and Luo X (2014), "Potential Mechanisms of Carbon Monoxide and High Oxygen Packaging in Maintaining Color Stability of Different Bovine Muscles", *Meat Sci.*, Vol. 97, pp. 189-196.
- Mace S, Joffraud J J, Cardinal M, Malcheva M, Cornet J, Lalanne V, Chevalier F, Serot T, Pilet M F and Dousset X (2013), "Evaluation of the Spoilage Potential of Bacteria Isolated from Spoiled Raw Salmon (*Salmo salar*) Fillets Stored Under Modified Atmosphere Packaging", *Int. J. Food Microbiol.*, Vol. 160, No. 3, pp. 227-238.
- Macé S, Joffraud J J, Cardinal M, Malcheva M, Cornet J, Lalanne V, Chevalier F, Sérot T, Pilet M F and Dousset X (2013), "Evaluation of the Spoilage Potential of Bacteria Isolated from Spoiled Raw Salmon (*Salmo salar*) Fillets Stored Under Modified Atmosphere Packaging", *Int. J. Food Microbiol.*, Vol. 160, pp. 227-238.
- Masniyom P, Benjakul S and Visessanguan W (2004), "ATPase Activity Surface Hydrophobicity Sulfhydryl Content and Protein Degradation in Refrigerated Seabass Muscle in Modified Atmosphere Packaging", *J. Food Biochem.*, Vol. 28, pp. 43-60.
- Masniyom P, Benjakul S and Visessanguan W (2002), "Shelf-Life Extension of Refrigerated Seabass Slices Under Modified Atmosphere Packaging", *J. Sci. Food and Agric.*, Vol. 82, pp. 873-880.
- Masniyom P, Benjama O and Maneesri J (2013), "Effect of Modified Atmosphere and Vacuum Packaging on Quality Changes of Refrigerated Tilapia (*Oreochromis niloticus*) Fillets", *Int. Food Res. J.*, Vol. 20, pp. 1401-1408.
- Meredith H, Valdramidis V, Rotabakk B T, Sivertsvik M, McDowell D and Bolton D J (2014), "Effect of Different Modified Atmospheric Packaging (MAP) Gaseous Combinations on *Campylobacter* and the Shelf-Life of Chilled Poultry Fillets", *Food Microbiol.*, Vol. 44, pp. 196-203.
- Mohan C O, Ravishankar C N, Srinivasa Gopal T K and Kumar A K (2009), "Nucleotide Breakdown Products of Seer Fish (*Scomberomorus commerson*) Steaks Stored in O₂ Scavenger Packs During Chilled

- Storage”, *Innovation Food Sci. Emerg. Tech.*, Vol. 10, pp. 272-278.
- Mohan C O, Ravishankar C N and Srinivasa Gopal T K (2007), “Effect of O₂-Scavenger on the Shelf-Life of Catfish (*Pangosiusutchi*) Steaks During Chilled Storage”, *J. Sci. Food and Agric.*, Vol. 88, pp. 442-448.
 - Noseda B, Islam M T, Eriksson M, Heyndrickx M, De Reu K, Van Langenhove H and Devlieghere F (2012), “Microbiological Spoilage of Vacuum and Modified Atmosphere Packaged Vietnamese Pangasiushypophthalmus Fillets”, *Food Microbiol.*, Vol. 30, pp. 408-419.
 - Ocano-Higuera V M, Marquez-Rios E, Canizales-Davila M, Castillo-Yanez F J, Pacheco-Aguilar R, Lugo-Sanchez M E, Garcia-Orozco K D and Graciano-Verdugo A Z (2009), “Postmortem Changes in Cazon Fish Muscle Stored on Ice”, *Food Chem.*, Vol. 116, pp. 933-938.
 - Pantazi D, Papavergou A, Pournis N, Kontominas M G and Savvaidis I N (2008), “Shelf-Life of Chilled Fresh Mediterranean Swordfish (*Xiphiasgladius*) Stored Under Various Packaging Conditions: Microbiological Biochemical and Sensory Attributes”, *Food Microbiol.*, Vol. 25, pp. 136-143.
 - Ravi-Sankar C N, Lalitha K V, Jose L, Manju S and Gopal T K S (2008), “Effect of Packaging Atmosphere on the Microbial Attributes of Pearlsport (*EtroplussuratensisBloch*) Stored at 0-2 °C”, *Food Microbiol.*, Vol. 25, pp. 518-528.
 - Restuccia D, Spizzirri G U, Parisi I O, Cirillo G, Curcio M, Iemma F, Puoci F, Vinci G and Picci N (2010), “New EU Regulation Aspects and Global Market of Active and Intelligent Packaging for Food Industry Applications”, *Food Control*, Vol. 21, pp. 1425-1435.
 - Seydim A C, Acton J C, Hall M A and Dawson P L (2006), “Effects of Packaging Atmospheres on Shelf-Life Quality of Ground Ostrich Meat”, *Meat Sci.*, Vol. 73, pp. 503-510.
 - Soccol M C H, Oetterer M, Cláudio Rosa Gallo C R, Marta Helena Fillet Spoto M H F and Denise Oliveira Biato D O (2005), “Effects of Modified Atmosphere and Vacuum on the Shelf Life of Tilapia (*Oreochromisniloticus*) Fillets”, *Braz. J. Food Technol.*, Vol. 8, pp. 7-15.
 - Thanonkaew A, Benjakul S, Vissessanguan W and Decker E A (2006), “The Effect of Metal Ions on Lipid Oxidation Color and Physicochemical Properties of Cuttlefish (*Sepia pharaonis*) Subjected to Multiple Freeze-Thaw Cycles”, *Food Chem.*, Vol. 95, pp. 591-599.
 - Thiansilakul Y, Benjakul S and Richards M P (2010), “Changes in Heme Proteins and Lipids Associated with Off-Odour of Seabass (*Latescalcarifer*) and Red Tilapia (*Oreochromismossambicus* × *Oniloticus*) During Iced Storage”, *Food Chem.*, Vol. 121, pp. 1109-1119.
 - Torrieri E, Antimo P, Silvana C, Vincenzo C, Ilaria F, Giovanna A, Buonocore G and Masi P (2011), “Effect of Modified Atmosphere and Active Packaging on the Shelf-Life of Fresh Bluefin Tuna Fillets”, *J. Food Eng.*, Vol. 105, pp. 429-435.
 - Wicklund R A, Paulson D D, Tucker E M, Stetzer A J, DeSantos F, Rojas M, MacFarlane B J and Brewer M S (2006), “Effect of Carbon Monoxide and High Oxygen Modified Atmosphere Packaging and Phosphate Enhanced, Case-Ready Pork Chops”, *Meat Sci.*, Vol. 74, pp. 704-709.
 - Yassoralipour A, Jamilah B, Russly A R and Fatimah A B (2012), “Biogenic Amines Formation in Barramundi (*Latescalcarifer*) Fillets at 8 °C Kept in Modified Atmosphere Packaging with Varied CO₂ Concentration”, *LWT – Food Sci. Tech.*, Vol. 48, pp. 142-146.
 - Yesudhasan P, Lalitha K V, Srinivasa Gopal T K and Ravishankar C N (2014), “Retention of Shelf Life and Microbial Quality of Seer Fish Stored in Modified Atmosphere Packaging and Sodium Acetate Pretreatment, Food Packag”, *Shelf Life*, Vol. 1, pp. 123-130.
 - Young H, Anang D M and Tiwari B K (2014), “Shelf Life and Textural Properties of Cooked-Chilled Black Tiger Prawns (*Penaeusmonodon*) Stored in Vacuum Pack or Modified Atmospheric Packaging at 4 or 20 °C”, *Food Packag*, *Shelf Life*, Vol. 2, pp. 59-64.
 - Yuan H, Kim B, Bødker S and Rosenvold K (xxxx), “Influence of Lamb Age and High-Oxygen Modified Atmosphere Packaging on Protein Polymerization of Long-Term Aged Lamb Loins”, *Food Chem.*, Vol. 135, pp. 122-126.
 - Zogula F O, Polata A and Ozogul Y (2004), “The Effects

of Modified Atmosphere Packaging and Vacuum Packaging on Chemical Sensory and Microbiological

Changes of Sardines (*Sardinapilchardus*)”, *Food Chem.*, Vol. 85, pp. 49-57.

APPENDIX

Table 1: Selected Methods of MAP Used for Various Types of Fish/Fish Product		
Reference	Method of MAP Used	Name of the Fish/Fish Product Used in MAP
(Soccol <i>et al.</i> , 2005)	The samples were packed under 60% CO ₂ /40% O ₂ modified atmosphere at a 2:1 (gas/fish) proportion, that is, 1000 mL of gas mixture per 500 g of fish	Nile tilapia (<i>Oreochromis niloticus</i>)
(Lauzon <i>et al.</i> , 2002)	The effects of modified atmosphere (CO ₂ /N ₂ : 60/40) in bulk storage of both redfish and the fish fillets. The samples were stored on ice in the cooling room (0-2 °C) till sensory rejection	Redfish
(Mace <i>et al.</i> , 2013)	Salmon cubes was placed in 3 plastic trays, each containing ~250 g portions (one for each analysis date), and packed under modified atmosphere (50% CO ₂ and 50% N ₂) and a low gas permeability film (low density polyethylene). All batches of inoculated salmon cubes and the control were stored at 8 °C for 12 days	Salmon (<i>Salmosalar</i>) fillets
(Emborg <i>et al.</i> , 2005)	Fresh tuna samples were stored either in Vacuum-Packed (VP) or Modified Atmosphere-Packed (MAP) at 1.0 8C for 28 days of storage. For MAP, 60% CO ₂ /40% N ₂ or 40% CO ₂ /60% O ₂ were used.	Tuna (<i>Thunnusalbares</i>)
(Fernández <i>et al.</i> , 2010)	Pieces fish samples (approximately 30 g of each) were packed into bags using modified atmospheres (initial CO ₂ content from 30 to 100%, remainder nitrogen) at a gas:product ratio of approximately 3:1 using a Technovac vacuum packer. The samples were then stored at a range of temperatures between 0 -10 °C	Atlantic salmon (Norway and Chile)
(Hansan <i>et al.</i> , 2007)	The samples of the fish were packaged. Two to three cod loins (around 500 g) were placed on trays (expanded polystyrene) with built-in absorbent drip pads. Each tray was put into a vacuum bag measuring 25 × 32.5 cm × 115 cm. A gas mixture of 50.0% CO ₂ -45.0% N ₂ - 5% O ₂ was injected to modify the atmosphere and the bags were heat-sealed individually and stored at temperature of 1.5 °C or -1 °C). The shelf-life of cod loins was evaluated	fresh cod loins (<i>Gadusmorhua L.</i>)
(Masniyom <i>et al.</i> , 2013)	Fillets samples weighing approximately 90-100 g were placed in vacuum bag (15 cm × 25 cm) with gas permeability (O ₂ transmission rate of 46.6 cm ³ m ⁻² day ⁻¹ at 38 °C, 1 atm pressure). The samples were prepared in different three packaging systems, system (1) packaged in air; (2) packaged using vacuum and (3) packed under MAP using a mixture of 60% CO ₂ , 10% O ₂ and 30% N ₂ . The bags were sealed and the samples were stored at 4 °C for 18 days.	Tilapia (<i>Oreochromis niloticus</i>) fillets

APPENDIX (CONT.)

Table 2: Shelf Life Extension in Some Types of Fish/Fish Product Packaged Using Various Methods of MAP			
Fish Product Used in MAP	Method of MAP Used	Observations and Shelf Life Extension	Reference
Catfish (Pangasiushypophthalmus)	The use of CO ₂ in modified atmosphere packaged Pangasius fillets significantly prolonged the shelf life compared to air and vacuum packaged fillets. Combination of 50% CO ₂ with 50% O ₂ appeared had an additional inhibitory effect on the microbiological growth, most probably on the lactic acid bacteria	MAP 1: 50% CO ₂ -50% N ₂ and MAP 2: 50% CO ₂ -50% O ₂) during storage at 4 °C. The shelf life of the fillets packaged in air, vacuum, MAP 1 and MAP 2 was estimated to be 7, 10, 12 and 14 days respectively.	Noseda <i>et al.</i> (2012)
Cod fillets	The sterile muscle samples of about 20 gram were placed in sterile petri dishes and incubated at 0 °C. Six The samples were incubated in atmospheres of 0, 50 and 100% CO ₂ , the remaining gas being N ₂ . The samples were analyzed just after preparation, after 2 weeks (0% CO ₂), after 3 weeks (50% CO ₂) and after 3 months (100% CO ₂)	H ₂ S producing bacteria was inhibited as <i>S. putrefaciens</i> maximum concentration found in MAP fish products was very low. Compared to VP, a shelf-life extension of 6-7 days was obtained with 48% CO ₂ in MAP. With pure CO ₂ the shelf life was only extended by 2-3 days	Paw Dalgaard <i>et al.</i> (1993)
Atlantic horse mackerel (Trachurstrachurus) fillets	The samples of the fish was packaged in a modified atmosphere (48% CO ₂ , 50% N ₂ and 2% O ₂) and kept at refrigeration temperature (6 °C) for 7 days	The samples showed that <i>Arthrobacter</i> , <i>Chryseobacterium</i> , <i>Photobacterium</i> , and <i>Pseudoclavibacter</i> (44.5% of total) dominated the microbial composition of the fish at the beginning of storage. It also showed <i>Yersinia</i> , <i>Serratia</i> , and <i>Shewanella</i> dominated at the late spoilage stages (over 57.2% of the total). <i>Carnobacterium</i> was identified at the beginning and end of the storage period. A shelf life of 5 days was obtained	Alfaro and Hernande (2013)
sardine (Sardinapilchardus)	quality and safety parameters of sardines were investigated using Vacuum Packaging (VP), and modified atmosphere packaging (MAP) (60% CO ₂ : 40% N ₂) for up to 15 days at 4 °C	Bacteria grew most quickly in sardine stored in air, followed by those in VP and the lowest counts were with MAP. For a period of 15 days, concentration of histamine, trimethylamine (TMA) increased to 20 mg/100 g, 13 mg/100 g and 10 mg/100 g for fish stored in air, VP and MAP, respectively. Shelf life extension was found to be 3 days in air, 9 days in VP and 12 days in MAP	Zogula <i>et al.</i> (2004)

