

**INTERNATIONAL JOURNAL OF FOOD AND
NUTRITIONAL SCIENCES**

IMPACT FACTOR ~ 1.021



Official Journal of IIFANS

SUSTAINABLE PREVENTION OF RENAL CALCULUS IN URINARY SYSTEM

Barbara Pyszczyk^{1*}

*Corresponding Author: Barbara Pyszczyk, ✉ basia.py@gmail.com

Received on: 13th March, 2017Accepted on: 7th August, 2017

Introduction: Kidney stone (renal calculus, lat. nephrolithiasis, urolithiasis) is the third most frequent urinary disease. It occurs on the entire world and affects up to 5% of the population between 30 and 50 years old which indicates that the disease is a serious social issue. A significant role in etiology of the disease is a diet. **Aim:** The prevention of kidney stones in urinary system and their recurrence. Dietary recommendations in kidney stone may vary depending on the kind of kidney stones. Proper nutrition in people suffering from renal calculus may be crucial in treatment. The diet ought to cover an individual's calorie requirement and provide proper and balanced amount of macro-and micro-nutrients. The patient has to be provided with carefully elaborated diet plan according to their individual needs regarding renal function, age, nutritional condition and metabolic disorders. Since there is no way to analyze the precipitation, making particular nutrition recommendations is not always possible thus, there have been elaborated nutrition guidelines to prevent kidney stones. **Conclusion:** The most effective way of prevention of renal calculus is limiting the intake of animal proteins, increasing the consumption of fruit and vegetables, decreasing the amount of salt in diet, regulated intake of calcium and proper hydration of the system.

Keywords: Renal calculus, Kidney stones, Nutritional therapy, Prevention, Hypercalciuria

INTRODUCTION

Kretowicz and Mianitius (2005) describe kidney stones (lat. *nephrolithiasis*, *urolithiasis*) as a chronic, systemic disease that is characterized by creating and existence of calculi in renal pelvis and kidney parenchyma. The calculi is created by components of either healthy or pathologically changed urine. The disease may be primary (the causes of metabolic orders are unknown) or secondary (e.g., urinary system defect, small intestine diseases, disturbances in the serum calcium and phosphorus concentration, caused by other systemic diseases).

Calculi is precipitated in urinary tract as a result of disturbances between crystalloids and natural crystallization inhibitors in urine. The factor catalyzing this process is

bacteria and lining, inducing nucleation and the reaction of the urine facilitating precipitation of some of its compounds. When calcium salts are deposited in urinary tract, an individual suffers from calcium oxalate stones (Wieczorkiewicz-Plaza, 2008).

Kidney stones affect around 5% of adult population. Moreover, as much as 12% of population developed the disease in USA (Tiselius, 2003). According to National Health and Nutrition Survey the frequency of stones is increasing. In 1994 in United States stones were diagnosed in only 6,3% of men and 4,1% of women. In 2012 the number of cases of kidney stone raised to 10,6% in men population and 7,1% in women (Scales, 2012). The occurrence of the disease is conditioned by geographical region, race, sex

¹ Dissertation Written Under Supervision of Prof. Dr Danuta Rosolowska, Huszcz at Warsaw University of Life Sciences, SGGW, Faculty of Nutrition and Consumer Sciences.

and age (Julka, 2010). Women suffer from kidney stones 3-4 times less frequently than men (Herold, 2008). In spite of the fact that the disease is more frequent in men, the ratio of men affected by kidney stones to women has decreased from 3,4 to 1,3 (Strope, 2010). The highest prevalence rate was seen in those at age of 50 (Baker, 1993).

DIAGNOSIS AND TREATMENT

Urinary Stones are diagnosed through taking detailed medical history, physical examination, laboratory tests and diagnostic imaging (Bar and Starownik, 2006; and Pearle, 2014). Special consideration is given to eating habits and long term medicines-therapy, especially medicines affecting the reaction of urine such as ephedrine, calcium supplements, vitamins and drugs increasing the risk of stones such as indinavir, triamterene and carbonic anhydrase inhibitors (Mysliwiec and Brzóska, 2005).

Routine general examination and urine culture ought to be carried out. Analyzing the sediment enables identification of crystals and define the chemical composition of the calculi. In case of genetic load and recurrent kidney stones the urine to be tested should be taken from the period of 24 hours prior to the culture test. It is generally recommended to take urine samples twice. On the basis of the daily urine sample diuresis is defined. Urine is determined in terms of pH, excretion of calcium, phosphorus, uric acid, oxalates, trisodium citrate and creatinine (Parks, 1996 and 2002).

Blood is tested to determine calcium concentration, nonorganic phosphate minerals, magnesium, potassium, uric acid, urea and creatinine (Parks, 1996 and 2002). Increased level of uric acid indicates abnormality in purines metabolism. Determining the concentration of calcium in serum helps to diagnose possible primary hyperparathyroidism and hiperkalcemia (Parks, 1996).

When diagnosing kidney stone the crucial role is played by medical imaging (Bochniewicka, 2006). Spiral computed tomography without gadopentetic acid allows detecting small radiolucent urinary calculi invisible in radiography of abdominal cavity. Intravenous urography helps to diagnose small radiolucent calculi that may cause just insignificant urinary retention. Radiography of abdominal cavity is useful and recommended in recognizing calcium stones however, it is not efficient in detecting radiolucent stones. Ultrasonography test should be routinely conducted along with radiography as it allows

to estimate the degree of urinary retention, the condition of renal parenchyma and diagnosing the presence of radiolucent stones. Intravenous pyelogram is carried out in case of lack of renal function during urography test and hydronephrosis diagnosed on the basis of ultrasonography test of abdominal cavity (Bar, 2004).

Not only does the choice of optimal treatment and preventing recurrence of the disease depend on the kind of kidney stone but also on proximal tubule transport disturbances (Gadomska-Prokop, 2007). Nutrients contributing to creation of kidney stones are first of all calcium, oxalates, sodium, magnesium, proteins and liquid (Haewook, 2015).

Elaborating dietotherapy aimed at treating kidney stones it is crucial to consider chemical composition of stones in urinary tract and factors we can influence by changing eating habits. These changes will considerably affect composition of urine and as a result will help to prevent recurrence of the disease.

PREVENTION

Acidity of urine as one of lithogenic factors is an important variable that is apparently affected by diet. The key factor for creating urinary calculi is not high excretion of uric acid but low urine pH level below 5,5. Treatment and prophylaxis of this type of stones consists in alkalization of urine to pH 6 level. This result can be obtained by increasing consumption of fruit and vegetables and reducing consumption of sources purines metabolized in system to uric acid. To avoid excessive alkalization that may lead to creation of uric acid crystals which may participate in heterogenic crystallization of calcium oxalate or tricalcium phosphate (Grases, 2006).

Tricalcium phosphate is a compound which is precipitated in alkaline urine and dissolved in acidic. Monitoring pH level is aimed at preventing precipitation of phosphate and dissolving calculi. The priority recommendation is to maximize dilution of urine through increased intake of fluids. Foods containing phosphates have to be eliminated from the diet. It is necessary to reduce consumption of legumes, alkaline mineral waters, potatoes, vegetables, fruit and cheeses. The protein consumption should be equivalent to about 0,8-1 g/kg mc/d (Wahl and Hess, 2000).

The detailed dietary guidelines related to lithogenic factor and type of kidney stone are presented in Table 1.

Table 1: Relation Between Lithogenic Factors in Urine, Type of Kidney Stone and Dietary Recommendations (Grases, 2006)

Lithogenic Factor	Biochemical Parameters of Urine	Kind of Stone	Diet Recommendations
Ph	<5,5	COM,	Reduce consumption of:
		UA,) Animal protein
		COM/UA,	Increase intake of:
		CYS) Citrus fruit juices
) Soft drinks
) Drinks rich in citric acid	
pH	>6	COM,	Reduce:
		COD,) Vegetarian diet
		HAP,) Fruit juices
		COD/HAP,) Soft drinks
		BRU) Drinks rich in citric acid
Calcium	Women	COD,	Increase consumption of:
	>250 mg/d	HAP,) Daily fluid intake (> 2 l/d)
	Men	COD/HAP) Reduce intake of:
	>300 mg/d) sodium
) animal protein
) vitamin C? (2 g/d)
Oxalates	>40 mg/d	COM	Reduce intake of:
) foods rich in oxalates
) vitamin C (2g/d)	
Citrates	<350 mg/d	COP,	Increase consumption of:
		COM) foods rich in citrates and drinks rich in citric acid
		COD,	
		HAP	
Phytins	<1 mg/d	COM,	Increase consumption of:
		COD,) foods rich in phytin
		BRU	
Urates	Women	UA,	Reduce consumption of:
	<600 mg/d	COM/UA) foods rich in
	Men) alcohol drinks
	>800 mg/d		

Note: COM – oxalic acid, COD – oxalic acid C₂H₂O₄, COD/HAP – oxalic acid C₂H₂O₄/hydroksyapatyt, HAP – hydroksyapatyt, STR – struvite, BRU – brushite, UA – uric acid, COM/UA – uric acid/calcium oxalate.

Since it is not always possible to enunciate particular dietary guidelines as there is not always a possibility to analyze the composition of urine, general dietary recommendations have been elaborated to prevent kidney stones.

Accurate recommendations supporting treatment of stones were described by Wahl and Hess (2000) in a so called reasonable diet. According to the authors the key aspect of prophylaxis concerning all types of stones is increase of fluid intake up to 3-4 liters a day which reduces supersaturation of urine (Cheungpasitporn, 2016). As diuresis should be maintained above the level of 2 l/d, it could be assumed that monitoring of excreting urine is more important than monitoring amounts of fluids intake. Liquids ought to be consumed in small amounts through the day after meals and before going to sleep. The intake of fluids should be increased in case of heat and physical effort. It has been proven that consuming less than 1 liter of fluids and producing less than 1 liter of urine per day contributes to creation of calcium stones. Eight-year prospective study conducted by Curhana (2004) proved that the risk of kidney Stones is 32% lower in people consuming 2700 ml of fluids per day than in people consuming less than 1400 ml of fluids per day.

Apart from amount of fluids consumed their quality is an important factor for the risk of lithogenicity. It is highly recommended to drink hypotonic solutions moderately enriched in calcium and magnesium. Wine, beer, coffee, tea and orange juice reduce the risk of calcium stones whereas grapefruit, apple and tomato juices increase the risk (Keble, 2004).

Coffee, in spite of containing oxalate, can be qualified as a drink reducing the risk of stones due to effect of caffeine that increases diuresis. It was found that 240 ml of coffee per day reduces the risk of deposits of calculi by 10% and a glass of tea by 8-14% (Curhan, 1996). Herbal and fruit infusions also limit the risk of stones (Itoh, 2005). A glass of grapefruit juice a day increases the risk of stones even by up to 40% (Curhan, 1998). According to the study 240 ml of beer or wine in everyday diet reduces the risk of stones approximately by 12% in women and 21% in men (Talalaj, 2009). There are no clear results concerning influence of consuming fizzy drinks (Curhan, 1998) and coke drinks (Curhan, 1996; and Passman, 2009).

Meals containing animal protein should be consumed 5-7 times a week but not twice a day. It is recommended to

reduce intake of animal proteins to 0,8-1,0 g per kg body weight, which allows reducing calciuria and excreting more citrates with urine (Talalaj, 2009), or replacing it with plant protein.

Reducing consumption of animal protein should be focused on eliminating from diet products containing big amounts of purines (jus, bones residue, giblets, game, lamb, sardines, herring, sprat).

Calcium in diet reduces the risk of calcium stones. In digestive tract it bonds oxalates and inhibits their absorption which results in reducing their concentration in blood and excreting with urine (Brzóska and Myœliwiec, 2009; and Taylor, 2013). Thus, a diet low in calcium is a risk of stones factor, in particular oxalate stones. It was found that consuming not enough amount of calcium contributes to absorption of oxalate and it's excreting with urine. The results of research conducted by Marengo and Romani (2008) indicate that if diet consists of at least 4 g of calcium per day, then even significantly increased amount of oxalate in consumed food does not increase its excreting. People consuming 1000-1300mg of basic calcium per day are the group of the least risk of stones in urine tract (Borghi, 2002; and Stamatelou, 2003).

Thus, it can be concluded that in dietotherapy of kidney stones calcium should not be reduced. Negative calcium balance leads to demineralization of skeletal system, osteoporosis and arterial hypertension. Limiting calcium intake in diet induces secondary hyperoxaluria.

Fruit and vegetables should be consumed at least once a day. However, products with high bioavailability of oxalates such as spinach, rhubarb, beetroot, nuts, chocolate, tea, wheat bran and strawberries should be eliminated as they cause significant increase of concentration of those compounds in urine. Recommended reduction of consumption of foods rich in oxalates is based on conclusion that calcium oxalate is a key factor for creating kidney stones. Reducing consumption of calcium increases absorption of oxalate and its excreting through kidneys whereas higher intake may decrease absorption of oxalates through binding their bigger amounts in intestines. Due to calcium's inhibiting influence on excreting oxalate with urine it is recommended to follow a diet aimed at regulation of calcium balance. The recommended sources of calcium are milk, yoghurt and kefir (Unruh, 2004; and Patchae, 2009).

The opposite effect is induced by a diet low in calcium, where reduced amount of calcium causes increase of

concentration of free ions of oxalate in intestines to be absorbed. As a result of excessive excretion of oxalates in urine (hyperoxaluria) (Unruh, 2004; and Patchae, 2009) calcium-oxalate stones may develop, which is a typical uric tract stones in children and adults (Lewandowski and Rodgers, 2004).

Higher consumption of fruit and vegetables resulted in increased intake of fibre, potassium and magnesium. Higher intake of potassium in diet reduces the risk of stones. Potassium prevents development of hyperglycemia that stimulates urethral reabsorption of citrates, substantial inhibitors of crystallization, reducing their concentration in urine (Brzósko and Myćliwiec, 2009). Magnesium, on the other hand, is the inhibitor of increase of calcium phosphate crystals (Rózański, 2009). The next recommendation for prevention of stones is reducing intake of sucrose and salt. Borghii (2002), Meschi (2004) and Nouvenne (2010) found that excessive intake of sodium chlorium contributes to development of lithogenic processes and proved that high content of sodium in food increases the amount of calcium in urine and every modification of the diet in order to decrease sodium and animal protein intake inhibits developing kidney stones.

Excessive intake of vitamin C, which is metabolized to oxalates, increases concentration of oxalates in urine (Traxer, 2003). Taylor (2004) recommend daily intake of vitamin C below 2 g.

Dietotherapy recommendations were updated by Tiselius (2008), Fink (2013) and Pearle and Skolarikos (2015). The new guidelines concerning treatment and prevention of kidney stone were based on findings concluded from randomized clinical studies and meta-analysis indicated in publications.

When the system does not react to diet modifications it is necessary to support dietotherapy with pharmacological treatment. Application of thiazides is of great importance as they decrease excreting of calcium and additionally has good influence on mineral density of bones (Heilberg, 1997; and Escribani, 2009). As the main side effect of intake of thiazides is hypokalemia that leads to lower excreting of citrates with urine, thiazides ought to be taken along with potassium citrate (Mike, 2006). Allopurinol is also significant in pharmacological treatment as it inhibits production of uric acid in urine.

In terms of prevention of urinary stones careful consideration must be given to natural therapy. Very good

results are obtained from drinking herbal infusions (among others Debelizyna, Urostonex, Urosept, Urosan, Rowatinex, Cystone, Fitolizyna, birch and nettle infusions, Uro-up, Zuravit, Furoxin, etc.) as they have anti-inflammatory, antibacterial, diuretic properties and help to relax smooth muscle of urinary tract which enhances excretion of calculi. Crenotherapy, therapy through intake of waters: Jan, Marysieńka, Józef, Wielka Pieniawa, which are supposed to mechanically scour small calculi, affects water and mineral balance and prevent infection of urinary tract.

SUMMARY

In spite of long-term studies on diet in pathogenesis of kidney stones there is no agreement between researchers on which of dietary factors directly contribute to recurrence of urinary stones (Prezioso, 2015). It can be concluded that the most efficient aspect of dietotherapy aimed at prevention of recurrence of urinary stones is reducing consumption of animal protein and salt, high intake of fruit and vegetables, normative intake of calcium and proper rehydration of the system.

ACKNOWLEDGMENT

I hereby represent that the work presented herein is original and does not infringe any copyrights, property rights or any related rights of third parties. The work has not been published so far. After publication the author waives any rights to the work presented.

REFERENCES

- Alatab S, Pourmand G, El Howairis Mel F, Buchholz N, Najafi I, Pourmand M R, Mashhadi R and Pourmand N (2016), "National Profiles of Urinary Calculi: A Comparison Between Developing and Developed Worlds", *Iran J Kidney Dis*, Vol. 10, pp. 51-61.
- Baker P W, Coyle P, Bais R and Rofe A M (1993), "Influence of Season, Age and Sex on Renal Stone Formation in South Australia", *Med J A.*, Vol. 159, pp. 390-392.
- Bar K and Starownik R (2006), "*Maloinwazyjne metody leczenia kamicy układu moczowego*", *Przegląd Urologiczny*, Vols. 7/6, pp. 24-26.
- Bar K (2004), "*Podręcznik urologii*", *Diagnostyka i leczenie*, Wydawnictwo Czelej, Lublin.
- Bochniewska V, Jung A, Giszczyk A, Muszyńska J and Krac E (2006), "*Kamica układu moczowego u dzieci* –

- inhibitory i promotory krystalizacji*”, *Pediatrics Medycyna Rodzinna*, Vol. 2, pp. 91-99.
- Borghi L, Schianchi T, Meschi T, Guerra A, Allegri F, Maggiore U and Novarini A (2002), “Comparison of Two Diets for the Prevention of Recurrent Stones in Idiopathic Hypercalciuria”, *N. Engl. J. Med.*, Vol. 346, pp. 77-84.
 - Brzósko S Z, Myœliwiec M [w:] and Myœliwiec M C (red.) (2009), “*Nefrologia, t. I. Wyd*”, pp. 320-329, Medical Tribune Polska, Warszawa.
 - Cheungpasitporn W, Rossetti S, Friend K, Erickson S B and Lieske J C (2016), “Treatment Effect, Adherence, and Safety of High Fluid Intake for the Prevention of Incident and Recurrent Kidney Stones: A Systematic Review and Meta-Analysis”, *J Nephrol.*, Vol. 29, pp. 211-219.
 - Curhan G C, Willett W C, Knight E L and Stampfer M J (2004), “Dietary Factors and the Risk of Incident Kidney Stones in Younger Women: Nurses’ Health Study II”, *Arch Intern Med*, Vol. 164, pp. 885-891.
 - Escribano J, Balaguer A, Roqué i, Figuls M, Feliu A and Ferre N (2014), “Dietary Interventions for Preventing Complications in Idiopathic Hypercalciuria”, *Cochrane Database Syst Rev.*, Vol. 11, p. 2.
 - Fink H A, Wilt T J, Eidman K E, Garimella P S, MacDonald R, Rutks I R, Brasure M, Kane R L, Ouellette J and Monga M (2013), “Medical Management to Prevent Recurrent Nephrolithiasis in Adults: A Systematic Review for an American College of Physicians Clinical Guideline”, *Ann Intern Med*, Vol. 158, pp. 535-543.
 - Gadomska-Prokop K (2007), “*Leczenie zachowawcze kamicy uk³adu moczowego*”, *Przegl³d Urologiczny*, Vol. 8, pp. 52-55.
 - Grases F, Costa-Bauza A and Prieto R M (2006), “Renal Lithiasis and Nutrition”, *Nutrition J*, p. 5.
 - Haewook Han, Adam M Segal, Julian L and Seifter Johanna T (2015), “Nutritional Management of Kidney Stones (Nephrolithiasis)”, *Clin Nutr Res.*, Vol. 4, pp. 137-152.
 - Heilberg I P, Martini L A, Szejnfeld V L and Schor N (1997), “Effect of Thiazide Diuretics on Bone Mass of Nephrolithiasis Patients with Idiopathic Hypercalciuria and Osteopenia”, *J Bone Miner Res*, Vol. 12, p. 479.
 - Herold G (2008), “*Medycyna wewnêtrzna*”, Tom 1-2, *Repetitorium dla studentów medycyny i lekarzy*, PZWL.
 - Itoh Y, Yasui T, Okada A, Tozawa K, Hayashi Y and Kohri K (2005), “Preventive Effects of Green Tea on Renal Stone Formation and the Role of Oxidative Stress in Nephrolithiasis”, *The Journal of Urology*, Vol. 173, pp. 271-275.
 - Jawalekar S, Surve V T and Bhutey A K (2010), “Twenty Four Hours Urine and Serum Biochemical Parameters in Patients with Urolithiasis”, *Nepal Med Coll J*, Vol. 12, pp. 5-7.
 - Julka S, Gupta S A and Srivastava A (2012), “Protocol Based Metabolic Evaluation in High Risk Patients with Renal Stones in North India”, *Indian Journal of Endocrinology and Metabolism*, Vol. 16, pp. 283-287.
 - Kretowicz M and Manitus J (2005), “*Kamica nerkowa a nadciœnienie têtnicze – wybrane aspekty*”, *Nadciœnienie têtnicze*, Vol. 5, pp. 327-334.
 - Kebler T, Jansen B and Hesse A (2002), “Effect of Blackcurrant-, Cranberry-, and Plum Juice Consumption on Risk Factors Associated with Kidney Stone Formation”, *Eur J Clin Nutr*, Vol. 56, pp. 1020-1023.
 - Marengo S R and Romani A M P (2008), “Oxalate in Renal Stone Disease: The Terminal Metabolite that Just Won’t Go Away”, *Nature Clinical Practice Nephrology*, Vol. 4, pp. 368-377.
 - Massey L K and Kynast-Gales S A (2001), “Diets with Either Beef or Plant Proteins Reduce Risk of Calcium Oxalate Precipitation in Patients with a History of Calcium Kidney Stones”, *J Am Diet Assoc*, Vol. 101, pp. 326-331.
 - Meschi T, Maggiore U, Fiaccadori E, Schianchi T, Bosi S, Adorni G, Ridolo E, Guerra A, Allegri F, Novarini A and Borghi L (2004), “The Effect of Fruits and Vegetables on Urinary Stone Risk Factors”, *Kidney Int*, Vol. 66, pp. 2402-2410.
 - Mike B, Siroky M B, Oates R D and Babayan R K (2006), “*Podrêcznik urologii*”, Wydawnictwo Czelej, Lublin.
 - Nouvenne A, Meschi T, Prati B, Guerra A, Allegri F, Vezzoli G, Soldati L, Gambaro G, Maggiore U and Borghi L (2010), “Effects of a Low-Salt Diet on Idiopathic Hypercalciuria in Calcium-Oxalate Stone Formers: A 3-

- mo Randomized Controlled Trial”, *Am J Clin Nutr*, Vol. 91, pp. 565-567.
- Nishiura J L, Martini L A, Mendonça C O, Schor N and Heilberg I P (2002), “Effect of Calcium Intake on Urinary Oxalate Excretion in Calcium Stone-Forming Patients”, *Braz J Med Biol Res*, Vol. 35, pp. 669-675.
 - Passman C M, Holmes R P, Knight J, Easter L, Pais V and Assimos D G (2009), “Effect of Soda Consumption on Urinary Stone Risk Parameters”, *J Endourol*, Vol. 23, pp. 347-350.
 - Parks J H and Coe P L (1996), “The Financial Effects of Kidney Stone Prevention”, *Kidney Int*, Vol. 50, pp. 1706-1712.
 - Pearle M S, Goldfarb D S, Assimos D G *et al.* (2014), “Medical Management of Kidney Stones: AUA Guideline”, *J Urol*, Vol. 192, pp. 316-324.
 - Prezioso D, Strazzullo P, Lotti T, Bianchi G, Borghi L, Caione P, Carini M, Caudarella R, Gambaro G, Gelosa M, Guttilla A, Illiano E, Martino M, Meschi T, Messa P, Miano R, Napodano G, Nouvenne A, Rendina D, Rocco F, Rosa M, Sanseverino R, Salerno A, Spatafora S, Tascia A, Ticinesi A, Travaglini F, Trinchieri A, Vespasiani G and Zattoni F (2015), “CLU Working Group: Dietary Treatment of Urinary Risk Factors for Renal Stone Formation: A Review of CLU Working Group”, *Arch Ital Urol Androl*, Vol. 7, No. 87, pp. 105-120.
 - Rózański W (2009), “Zapobieganie nawrotom kamicy wapniowej, Charakterystyka kamieni zawierających sole wapnia”, *Człowiec II, Przegląd Urologiczny*, Vol. 10, No. 6, pp. 34-36.
 - Scales C D, Smith A C, Hanley J M and Saigal C S (2012), “Prevalence of Kidney Stones in the United States”, *Urologic Diseases in America Project*, Vol. 62, pp. 160-165.
 - Skolarikos A, Straub M, Knoll T, Sarica K, Seitz C, Petrik A and Türk C (2015), “Metabolic Evaluation and Recurrence Prevention for Urinary Stone Patients: EAU Guidelines”, *Eur Urol*, Vol. 67, pp. 750-763.
 - Stamatelou K K, Francis M E, Jones C A and Curhan G C (2003), “Time Trends in Reported Prevalence of Kidney Stones in the United States: 1976-1994”, *Kidney International*, Vol. 63, pp. 1817-1823.
 - Stroppe S A, Wolf J S Jr and Hollenbeck B K (2010), “Changes in Gender Distribution of Urinary Stone Disease”, *Urology*, Vol. 75, pp. 543-546.
 - Swatowski A (2009), “Leczenie zachowawcze kamicy moczowej”, *Forum Nefrologiczne*, Vol. 2, No. 3, pp. 189-195.
 - Talalaj M and Tobola M (2009), “*Marcinkowska-Suchowierska E.: Kamica układu moczowego – prewencja i leczenie farmakologiczne*”, *Postępy Nauk Medycznych*, Vol. 5, pp. 387-394.
 - Taylor P L and Curhan G C (2013), “Dietary Calcium from Dairy and Nondairy Sources, and Risk of Symptomatic Kidney Stones”, *J Urol*, Vol. 190, pp. 1255-1259.
 - Taylor E N, Stampfer M J and Curhan G C (2004), “Dietary Factors and the Risk of Incident Kidney Stones in Men: New Insights After 14 Years of Follow-Up”, *Journal of the American Society of Nephrology*, Vol. 1, pp. 3225-3232.
 - Tiselius H G, Alken P, Buck C, Gallucci M, Knoll T, Sarica K and Türk C (2008), “Guidelines on Urolithiasis”, p. 128, Arnhem, The Netherlands: European Association of Urology (EAU).
 - Tiselius H G (2003), “Epidemiology and Medical Management of Stone Disease”, *British Journal of Urology International*, Vol. 91, pp. 758-767.
 - Tracy C R, Best S, Bagrodia A, Poindexter J R, Adams-Huet B, Sakhae K, Maalouf N, Pak C Y and Pearle M S (2014), “Animal Protein and the Risk of Kidney Stones: A Comparative Metabolic Study of Animal Protein Sources”, *J Urol*, Vol. 192, pp. 137-141.
 - Traxer O, Pearle M S, Gattegno B and Thibault P (2003), “Vitamin C and Stone Risk: Review of the Literature”, *Prog Urol*, Vol. 13, pp. 1290-1294.
 - Von Unruh G E, Voss S, Sauerbruch T and Hesse A (2004), “Dependence of Oxalate Absorption on the Daily Calcium Intake”, *J Am Soc Nephrol*, Vol. 15, pp. 1567-1573.
 - Wahl Ch and Hess B (2000), “Nierensteine – ist die Ernährung Trigger oder Therapeutikum?”, *Ther Umschau*, Vol. 57, pp. 138-145.
 - Wiczkiewicz-Plaza A, Sikora P, Bieniasz B, Borzêcka H, Plaza P and Zajaczkowska M (2008), “Leczenie pierwszych epizodów kamicy u dzieci w materiale własnym”, *Polski Merkuriusz Lekarski*, Vol. XXIV, Supplement 4, pp. 68-70.



IJEANS

**International Journal of Food
And Nutritional Sciences**