

**INTERNATIONAL JOURNAL OF FOOD AND
NUTRITIONAL SCIENCES**

IMPACT FACTOR ~ 1.021



Official Journal of IIFANS

PROXIMATE COMPOSITION, FUNCTIONAL PROPERTIES AND
ACCEPTABILITY OF WHEAT BASED FUNKASO AS AFFECTED BY ADDITION OF
PEARL MILLET AND SOYBEAN FLOURS

Fannah Mustapha Adam¹, Mamudu Halidu Badau^{1*}, Amin Odera Igwegbe¹
and Hadiza Kubura Lawan¹

*Corresponding Author: Mamudu Halidu Badau, ✉ badau@unimaid.edu.ng

Received on: 22nd April, 2018

Accepted on: 14th August, 2018

Flours of wheat cultivars, pearl millet and soybean were used to formulate funkaso blends. The experimental design was a 3 x 4 x 2 factorial experiment. Factor one was wheat cultivar (Norman, Cettia, Atilla gan Atilla), factor two the substitution level (0, 20, 30, 40%) of wheat cultivars for pearl millet flour and factor three substitution level (0, 30%) of cereal (combined wheat and pearl millet flours) for soybean flour giving rise to 24 samples plus one commercial sample making a total of 25 samples all together. Funkaso was produced from the 25 blends and their proximate composition, functional properties and acceptability determined. Data generated were subjected to statistical analysis. There were wide variations among the 25 funkaso formulations with regard to proximate composition, functional properties and sensory score by panelists. Addition of soybean increased the protein and fat contents of complementary food significantly ($P < 0.05$). Sensory scores tasted by panelists were generally high and therefore wheat can be supplemented with soybean and pearl millet at 30% and 28% levels, respectively in funkaso processing which can increase protein and fat contents as well as increase the profit margin and save foreign exchange earnings by reducing importation of wheat.

Keywords: Funkaso, Soybean, Pearl millet, Wheat

INTRODUCTION

Funkaso is a prestigious well liked and consumed traditional fermented foods in northern Nigeria. It is of great importance in the diet of kanuris and shuwa Arab. It is prepared mostly on special occasions and festive periods because of its rich value. Method for preparation of *Funkaso* varies from one processing to other due to lack of standardized ingredient formulation that would ensure product consistency.

It is prepared by mixing of whole wheat flour (fine or grits) together with water, yeast, baking powder and pint of salt to form a batter (thin flour mixtures that are beaten or stirred), which is allowed to stand depending on weather

condition and then deep fat fried in oil. Wheat is the principal ingredient used for *funkaso* production but it remains costly despite the intensity in local production in Nigeria. Most of the wheat used in Nigeria is imported and the negative effect of this on foreign exchange cannot be over emphasized. This prompts investigation into supplementation of wheat with any readily available cereal grains.

Pearl millet is abundant in semi-arid region of northern Nigeria where *funkaso* is usually produced and consumed, and incidentally it has higher protein content compared to most cereal grains (Badau *et al.*, 2008; and Badau *et al.*, 2009). Therefore, there is need to supplement wheat with

¹ Department of Food Science and Technology, Faculty of Engineering, University of Maiduguri, P. M. B. 1069, Maiduguri, Nigeria.

pearl millet in *funkaso* production, if the industry is to thrive. Supplementing of wheat with pearl millet in *funkaso* production can only reduce the cost of production and probably increase its profit margin thereby saving our foreign exchange but there is the problem of incomplete essential amino acid profile, since both wheat and pearl millet are cereal grains.

The nutritional quality of most cereal protein is poor (Wang *et al.*, 2008) because they contain less of the essential amino acid particularly lysine needed for growth and maintenance. On the other hand legumes contain lysine but lack methionine and cystine essential amino acid abundant in cereals. Therefore, blending both cereal and legumes in *funkaso* production can make available the required essential amino acid needed by the body comparable to protein of meat, fish and eggs.

Soyabean (*Glycine max*) is a legume that is considered as an excellent source of protein (35-40%) and is not expensive. The seed is the richest in food value of all plant foods consumed in the world (Kure *et al.*, 1998). Soybean protein have been used widely in foods for their distinctive physico-chemical and functional properties as well as nutritional value (Wang *et al.*, 2008). Therefore, its utilization as a supplement in production of *funkaso* may not only address its nutrient status but also functional properties (reducing oil absorption) thereby making it more economical and affordable to the low income earners.

In a similar studies, Badau and Magaji (2011) and Badau *et al.* (2013) supplemented pearl millet with cowpea in alkaki and garabia (traditional wheat and rice based snack). Kalmajit *et al.* (2013)-incorporate cowpea flour as functional ingredients in wheat baked foods. Hence, incorporation of legume in *funkaso* (wheat based product) will go a long way in addressing its deficit. It has been reported that mutual compensation is closest to ideal when the ratio by mass of cereal to legume is roughly 70:30 (Marero *et al.*, 1988; and FAO, 2005).

The problems associated with the local production include: Non-standardization of equipment, process and raw materials, inadequate hygiene during and after production, and little or no packaging. These problems can result in poor preservation techniques and high levels of contaminant in food resulting in food borne illness (Ingbian and Akpapunam, 2005). These problems could be tackled by, among others, standardizing ingredients and processes in *funkaso* production.

The objectives of the study were to produce *funkaso* from the blend of wheat, pearl millet and soybean flours at various proportions; evaluate the flour blends and *funkaso* produced from various formulations by determining their functional properties (bulk density, water and oil absorption, swelling capacity), proximate composition and acceptability.

MATERIALS AND METHODS

Three wheat cultivars (*Norman*, *Cettia* (CTA) and *Atila* gan *Atila*), one Pearl millet cultivar (SOSAT C-88) and Soybean were obtained from Lake Chad Research Institute Maiduguri, while other ingredients (baking powder, yeast and oil) were obtained from Maiduguri Monday market, Nigeria.

Whole Wheat Flour Production

The wheat grain was cleaned to remove unwanted chaff and dirt; it was then milled into fine flour without conditioning so as to prevent separation of the bran from the endosperm which is not desired in production of whole wheat flour for *funkaso* production.

Pearl Millet Flour Production

The grain was cleaned to remove foreign matters, conditioned to soften the bran, mellow the endosperm hence facilitating its separation during dehulling. The dehulled grain is then washed with water and allowed to dry in the sun, milled in a hammer mill and finally sieved to obtain fine flour.

Soybean Flour Production

Soybean was sorted, washed and soaked for 5 hours in a clean water of three times its weight and volume until the coat becomes soaked and wet to enhance the removal of some soluble anti-nutrient factor and facilitate dehulling. The soybean was further washed, drained and partially sundried. The soybean was then toasted at surface temperature of 180 ± 5 °C for 30 minutes in an open thick aluminum pot (Iwe, 2003; and Badau *et al.*, 2006). It was milled into fine flour with a hammer mill and let to pass through a 0.8 μ m mesh size screen.

Standardization Procedure

Funkaso mix flour was standardized by paying several visits to traditional *funkaso* processors in Maiduguri following the procedure of Badau *et al.* (1997). A weighing balance was used to measure out all the ingredients added by the traditional processors during the preparation. The average weights of the various ingredients used in the traditional

Table 1: *Funkaso* Formulations

Formulations (Codes)	Whole Wheat Flour (g)	Pearl Millet Flour (g)	Soybean Flour (g)	Water (ml)	Baking Powder (g)	Yeast (g)	Salt (g)
CC	200	-	-	210	3	2.4	3.6
A1	200	-	-	210	3	2.4	3.6
A2	140	-	60	210	3	2.4	3.6
A3	160	40	-	210	3	2.4	3.6
A4	112	28	60	210	3	2.4	3.6
A5	140	60	-	210	3	2.4	3.6
A6	103	42	60	210	3	2.4	3.6
A7	120	80	-	210	3	2.4	3.6
A8	84	28	60	210	3	2.4	3.6
C1	200	-	-	210	3	2.4	3.6
C2	140	-	60	210	3	2.4	3.6
C3	160	40	-	210	3	2.4	3.6
C4	112	28	60	210	3	2.4	3.6
C5	140	60	-	210	3	2.4	3.6
C6	103	42	60	210	3	2.4	3.6
C7	120	80	-	210	3	2.4	3.6
C8	84	28	60	210	3	2.4	3.6
N1	200	-	-	210	3	2.4	3.6
N2	140	-	60	210	3	2.4	3.6
N3	160	40	-	210	3	2.4	3.6
N4	112	28	60	210	3	2.4	3.6
N5	140	60	-	210	3	2.4	3.6
N6	103	42	60	210	3	2.4	3.6
N7	120	80	-	210	3	2.4	3.6
N8	84	28	60	210	3	2.4	3.6

recipe was used to produce the as a basis for arriving at the formulations presented in Table 1.

Experimental Design

A $3 \times 4 \times 2$ factorial experimental design as reported by Gomez and Gomez (1983) was used for the production of

funkaso flour blends, where three (3) wheat cultivars {*Norman*, *Cettia* (CTA), *Atilla gan Atilla*} were substituted with pearl millet (SOSAT C-88) at four (4) levels (0, 20, 30, 40%) and soybean (ER-biu) at two (2) levels (0, 30%) giving rise to a total of 24 samples plus commercial one as control.

Formulation of Funkaso Flour Blends

The flour blend ratios used were 100%; 70%:00:30%; 80%:20%:0%; 56%:14%:30%; 70%:00%:30%; 49%:21%:30%; 60%; 40%:00%; and 42%:28%:30%. Other quantity of ingredients salt, yeast, baking powder and water remained the same for all formulations (Table 1).

Funkaso Production

Funkaso was prepared by mixing of whole wheat flour with together water, yeast, baking powder and pint of salt to form a batter. The batter was allowed to stand depending on weather condition (2-4 hrs) and then deep fried in oil (Figure 1).

Figure 1: Flow Diagram of Modified *Funkaso* Preparation



Functional Properties

Water absorption and Oil absorption capacities of the samples were determined using the method described by Ojinnaka *et al.* (2013). One gramme (1 g) of the flour was mixed with 10 ml of the water in a centrifuge tube and allowed to stand at room temperature for 1 hr. This was centrifuged at 3500 rpm for 30 min. The volume of water in the sediment was measured. The water absorption was calculated as water absorbed per gramme of flour (Adejuyitan *et al.*, 2009). Bulk density of the samples was determined using the method described by Omidiran *et al.* (2015) and calculated as weight of flour (g) divided by flour volume (cm³) (Okaka and Potter, 1979; and Adejuyitan *et al.*, 2009). Water swelling capacity of the sample was determined by the method described by Okaka and Potter (1977) and Adejuyitan *et al.* (2009).

Proximate Composition

Moisture, protein, fat, ash and crude fibre were determined by methods of AOAC (2003). Carbohydrate was calculated as described by Asma *et al.* (2006) and Energy was calculated using Atwater factor (FAO, 2002).

Sensory Evaluation

The sensory evaluation test was conducted by a team of semi-trained 15 panelists drawn from staff and students of

University of Maiduguri. The sample were rated for taste, color, texture and overall acceptability based on nine point hedonic scale with representing 9 like extremely and 1 representing dislike extremely as described by Ihekoronye and Ngoddy (1985). Although, the panelists were selected based on basic requirements of a panelist, such as availability for the entire period of evaluation, interest, willing to serve, good health (not suffering from colds), not allergic or sensitive to the products evaluated (Penfield and Campbell, 1990).

Statistical Analysis

The data generated from the study were subjected to analysis of variance (ANOVA) as described by Gomez and Gomez (1983) and means separated using Duncan Multiple Range Test as described by Duncan (1955).

RESULTS AND DISCUSSION

Proximate Composition of Flours of Wheat Cultivars, Pearl Millet Cultivar, Soybean

The proximate composition of three wheat cultivars, one pearl millet cultivar and soybean are shown in Table 2. There were wide significant ($P < 0.05$) variations of moisture, protein, fat, ash, crude fibre, soluble carbohydrate and energy

Table 2: Proximate Composition of Wheat, Pearl Millet and Soybean

S. No.	Commodity	Proximate Composition (%)						Energy (Kcal/100 g)
		Moisture	Protein	Fat	Ash	Crude Fibre	Carbohydrate	
1	Wheat cultivars							
	Atilagan Atilla	11.10±0.00 ^d (88.90)	9.59±0.00 ^a	3.16±0.10 ^a	3.37±0.20 ^d	1.58±0.10 ^b	71.19±0.98 ^b ^c	351.58±0.47 ^a
	Cettia (CTA)	10.56±0.48 ^d (89.44)	11.65±0.00 ^b	3.09±0.18 ^a	3.24±0.00 ^c	1.57±0.00 ^b	69.88±0.49 ^b	353.97±1.82 ^{ab}
	Norman	9.42±0.33 ^c (90.58)	9.38±0.04 ^a	3.69±0.00 ^b	2.99±0.00 ^b	1.64±0.01 ^b	72.86±0.02 ^{cd}	362.23±0.15 ^b
2	Pearl millet cultivar							
	SOSAT C-88	7.49±0.11 ^b (92.58)	11.63±0.19 ^b	4.58±0.14 ^c	1.97±0.03 ^a	1.32±0.01 ^a	73.83±1.40 ^d	375.62±6.27 ^c
3	Soybean	4.16±0.00 ^a (95.84)	44.15±0.15 ^c	18.92±0.14 ^d	3.24±0.00 ^c	4.89±0.00 ^c	24.63±0.01 ^a	448.87±2.81 ^d

Note: Means (±SE) within each column not followed by the same superscripts are significantly different ($P < 0.05$). Values in brackets are percent dry matter beneath the moisture content of each commodity.

contents among the ingredients used for *funkaso* formulations. Highest moisture content was recorded by wheat cultivars (Cettia, Atilla gan Atilla and Norman) while soybean had the lowest ($P<0.05$). On the other hand, soybean had the highest ($P<0.05$) protein and fat contents. Norman had the highest ($P<0.05$) ash content. Atilla gan atilla had the highest crude fibre followed by Cettia and soybean with insignificant variation ($P<0.05$). The carbohydrate content of Norman and SOSAT C-88 were higher ($P<0.05$) than the rest of the ingredients. Soybean had the lowest ($P<0.05$) carbohydrate but had the highest ($P<0.05$) energy values.

Proximate Composition of *Funkaso* Flour Blends

The proximate composition of *funkaso* flour blends is presented in Table 3. It revealed that moisture being important criteria of food shelf stability and was found to range from 8.93-11.25%. The results showed that protein level increased significantly ($p<0.05$), ranging from 6.97-18.67%. Sample C2 had the highest value and C3 was observed to have the least value. Similar results were reported by Okoye *et al.* (2010) and Yusufu *et al.* (2013).

The fat content ranged from 1.88-8.81. Least value was observed in sample N7 (60% whole wheat flour, 40% pearl millet), which was significantly ($P<0.05$) different from N2 (70% whole wheat flour, 30% soybean) and C2 (70% whole wheat flour, 30% soybean). *Funkaso* flour blends (produced from A variety) were having higher amount of crude fat but significantly different from the control having a value of 14.17 ± 0.10 . An increase in the level of fat was observed with soybean incorporation, similar trend was observed by Ghangale and Jadhao (2016) in puri (Indian snack similar but produced from white flour) which was also supplemented with soybean up to 15%. These implies that *funkaso* from composite flour can be referred to as healthy and energy dense food because of high amount of protein, crude fibre and fat, hence will be recommended for those that are on strenuous work especially athletes and growing children as the case may be. Sample C1 (100% Whole wheat flour) and C3 (80% whole wheat flour, 20% pearl millet) remained insignificant ($P>0.05$) from control (CC).

Dietary fiber is one of the important nutraceutical components with wide range of health benefits. Pearl millet is one of the richest sources of iron and dietary fiber (Singh and Sehgal, 2008; and Gangale and Jadhao, 2016). The control varied significantly from all the sample with the

exception of sample N2 (70% whole wheat flour, 30% soybean) and N4 (56% whole wheat flour, 14% pearl millet, 30% soybean). The crude fibre content of *funkaso* produced ranged between 2.18-3.78. *Funkaso* being a whole wheat product is expected to have high amount of fibre. Sample supplemented with pearl millet and soybean had its level increased significantly at $p<0.05$ these could be as a result of relatively high amount of fibre in the bran of the whole-wheat flour and/or residual hulls from the supplements that have been inevitably included during processing, representing variable fraction of dietary fibre including mostly the lignin, cellulose and hemicelluloses components (Mannay and Shadaksharaswamy, 2005). The increased fibre and the lower carbohydrate content of *funkaso* have several health benefits, as it will aid digestion, reduce constipation and reduce the risk of colon cancer often associated with products from refined grain flours (Slavin, 2005; and Elleuch *et al.*, 2011).

There was significant increase ($P<0.05$) in ash content of the flour blends with a range of 1.33-3.87% with incorporation of soybean flour substitution especially in the A variety formulation. Sample N3 had the least and C6 the highest value. All the samples are significantly different $p>0.05$. Ash content is indicative of the amount of minerals in any food sample. The increase in ash is indicative of high mineral content of pearl millet and soybean flour.

A significant decrease in carbohydrate was observed with the incorporation of pearl millet and soybean ranging from 46.53-58.43%. The concomitant decrease in carbohydrate and increase in protein content is due to supplementation with pearl millet and soybean since soybean flour is rich in proteins and fats (Igbabul *et al.*, 2012). All the samples were significantly different from the CC with exception of C5 (70% whole wheat flour, 30% pearl millet, 00% soybean). The highest value was observed in sample C5 and the least in C6 (49% whole wheat flour, 21% pearl millet 30% soybean). No significant difference were observed in these samples A3 (80% whole 20% Pearl millet), A4 (56% whole wheat flour, 14% pearl millet, 30% soybean), A6 (42% whole wheat flour, 28% pearl millet, 30% soybean), A8 (42% whole wheat flour, 28% pearl millet, 30% soybean) and N2 among others.

Energy value range for all the samples is between 380.96-411.92 kcal, the CC (393.78) varied from all the samples except A5 (394.25) at $p<0.05$ level of significance. The high energy value in sample A3 (411.92) could be attributed to the high

fat contents of which has been absorbed by the sample during frying which is directly proportional to the intrinsic and extrinsic factors which determines the end product. This is highly desired especially in famine and war-torn locations where the next meal is not easy to come by. High-energy foods tend to have a protective effect in the optimal utilization of other nutrients (Wardlaw, 2004).

Proximate Composition of Funkaso Produced from Several Formulations

Proximate composition of *funkaso* produced from several

formulations is presented in Table 4. The results showed that the dry matter content which is sum of the total solid in the flour ranged between 88.91-91.06%. The moisture content accounts for 8.93-11.25%; with N4 (56% whole wheat flour, 14% pearl millet, 30% soybean) having the least value and C1 (100% Whole wheat flour) has the highest percentage. The moisture content decreased with pearl millet and soybean flour supplementation, samples formulation (C4, C5 and C8), (C6, C7) and (N1, N2) from variety A and C are insignificant different ($p < 0.05$). All other samples varied

Table 3: Proximate Composition Funkaso Flour Blends

Formulations				Proximate Composition (%)							Energy (Kcal/100 g)
WT	ST	SB		Dry Matter	Moisture	Crude Protein	Crude Fat	Crude Fibre	Ash	CHO	
100AA	0	0	(A1)	88.89±0.33 ^{def}	11.11±0.00 ^{ij}	9.59±0.00 ^h	3.16±0.01 ^{cd}	3.37±0.02 ⁱ	1.58±0.10 ^a	71.19±0.09 ^j	351.58±0.47 ^a
70AA	0	30	(A2)	89.86±0.00 ^{efgh}	10.14±0.00 ^{def}	18.30±0.00 ⁿ	7.74±0.06 ^c	3.22±0.07 ^p	1.21±0.07 ^f	59.39±0.04 ^{bc}	372.49±0.47 ^d
80AA	20	0	(A3)	89.98±0.02 ^{abc}	11.01±0.02 ^{hij}	7.82±0.03 ^c	2.88±0.06 ^{bc}	2.83±0.00 ^e	1.72±0.00 ^{bc}	73.71±0.08 ^{klm}	352.12±0.37 ^a
56AA	14	30	(A4)	90.04±0.01 ^{fghi}	9.95±0.01 ^{def}	15.66±0.02 ^{km}	6.62±0.06 ^{fg}	4.11±0.00 ^o	2.15±0.02 ^{def}	61.48±0.07 ^e	378.26±0.38 ^{def}
70AA	30	0	(A5)	89.13±0.00 ^{abcd}	10.86±0.00 ^{ghij}	10.52±0.00 ^j	2.70±0.01 ^{bc}	2.70±0.00 ^c	1.72±0.01 ^{bc}	71.55±0.03 ⁱ	352.73±0.14 ^a
49AA	21	30	(A6)	89.44±0.01 ^{bcde}	10.56±0.01 ^{fghi}	16.18±0.01 ^{km}	7.30±0.00 ^{efg}	3.88±0.00 ^l	2.19±0.00 ^{ef}	59.88±0.02 ^c	374.97±0.02 ^{de}
60AA	40	0	(A7)	89.43±0.38 ^{bcde}	10.56±0.38 ^{fghi}	9.61±0.01 ^h	2.33±0.00 ^{ab}	2.42±0.00 ^a	1.76±0.00 ^c	73.29±0.36 ^{jk}	352.65±1.59 ^a
42AA	28	30	(A8)	89.83±0.00 ^{efg}	10.16±0.00 ^{ef}	17.87±0.00 ^{mn}	6.98±0.00 ^{ef}	3.72±0.00 ^k	2.22±0.00 ^f	61.02±0.01 ^{ab}	375.46±0.07 ^{de}
100C	0	0	(C1)	89.44±0.48 ^{bcde}	10.56±0.48 ^{fghi}	11.65±0.00 ^k	3.09±0.01 ^c	3.24±0.00 ^h	1.57±0.00 ^a	69.88±0.49 ^b	353.97±1.82 ^{ab}
70C	0	30	(C2)	89.70±0.16 ^{def}	10.29±0.16 ^{fg}	18.67±0.01 ⁿ	8.48±0.19 ^g	4.02±0.00 ⁿ	2.08±0.00 ^d	67.44±0.33 ^g	376.78±0.35 ^{def}
80C	20	0	(C3)	88.91±0.04 ^{ab}	11.25±0.20 ^j	6.97±0.01 ^a	3.08±0.04 ^c	3.11±0.00 ^g	1.63±0.00 ^{ab}	73.94±0.16 ^{lm}	351.41±1.06 ^a
56C	14	30	(C4)	89.57±0.21 ^{def}	10.42±0.21 ^{fgh}	17.66±0.00 ^c	8.67±0.01 ^g	3.97±0.00 ^m	2.13±0.00 ^{def}	66.13±0.19 ^f	377.39±0.86 ^{def}
70C	30	0	(C5)	89.58±0.23 ^{def}	10.41±0.23 ^{fgh}	7.77±0.00 ^c	2.89±0.01 ^{bc}	2.95±0.01 ^f	1.70±0.00 ^{bc}	74.26±0.22 ^m	361.27±8.07 ^c
49C	21	30	(C6)	89.48±0.11 ^{bcdef}	10.51±0.11 ^{fghi}	17.13±0.08 ^{mn}	8.56±0.00 ^{fg}	3.85±0.03 ^l	2.16±0.00 ^{def}	61.78±0.17 ^c	376.69±0.57 ^{def}
60C	40	0	(C7)	89.85±0.07 ^a	10.81±0.40 ^{ghij}	8.81±0.00 ^f	8.81±0.00 ^{bc}	2.55±0.00 ^b	1.74±0.00 ^{abc}	73.38±0.40 ^{ijkl}	353.14±1.61 ^{ab}
42C	28	30	(C8)	89.52±0.04 ^{cdef}	10.48±0.04 ^{fgh}	17.64±0.00 ^{mn}	14.64±0.00 ^{efg}	3.66±0.05 ^j	2.17±0.00 ^{def}	60.70±0.04 ^a	376.41±0.13 ^{def}
100N	0	0	(N1)	90.57±0.03 ^{ijkl}	9.42±0.03 ^{abcd}	9.38±0.04 ^g	3.69±0.00 ^d	2.99±0.00 ^f	1.64±0.01 ^{ab}	72.86±0.02 ^l	362.23±0.15 ^c
70N	0	30	(N2)	90.49±0.28 ^{hijk}	9.51±0.28 ^{abcd}	18.37±0.01 ⁿ	8.26±0.02 ^{efg}	4.10±0.00 ^o	2.10±0.00 ^{de}	67.64±0.33 ^g	378.43±1.06 ^{def}
80N	20	0	(N3)	90.66±0.0 ^{ijkl}	9.33±0.00 ^{abc}	12.13±0.00 ^l	3.01±0.00 ^c	2.73±0.00 ^d	1.70±0.00 ^{bc}	71.08±0.01 ⁱ	359.95±0.09 ^{bc}
56N	14	30	(N4)	91.06±0.018	8.93±0.08 ^a	16.68±0.01 ^r	8.68±0.00 ^g	3.88±0.00 ^l	2.23±0.04 ^f	60.59±0.05 ^a	383.26±0.16 ^f
70N	30	0	(N5)	90.97±0.01 ^{kl}	9.02±0.01 ^{ab}	10.40±0.00 ^j	2.98±0.00 ^c	2.54±0.00 ^b	2.12±0.07 ^{def}	73.00±0.07 ^j	354.53±8.86 ^{ab}
49N	21	30	(N6)	90.40±0.00 ^{hij}	9.59±0.00 ^{bcde}	16.11±0.00 ^m	8.64±0.01 ^{fg}	3.71±0.00 ^k	2.16±0.00 ^{def}	62.77±0.00 ^d	381.28±0.10 ^{ef}
60N	40	0	(N7)	90.31±0.02 ^{ghi}	9.69±0.02 ^{cde}	7.69±0.00 ^b	1.88±0.94 ^a	2.51±0.00 ^b	1.72±0.02 ^{bc}	75.55±0.02 ^b	358.45±0.18 ^{abc}
42N	28	30	(N8)	90.44±0.00 ^{hijk}	9.55±0.00 ^{bcd}	17.17±0.00 ^{mn}	8.28±0.00 ^{efg}	3.64±0.00 ^j	2.32±0.00 ^g	61.02±0.00 ^{ab}	379.36±0.04 ^{def}

Note: Means (±SE) in the same column having different superscripts are significantly ($p < 0.05$) different. WT = Wheat; ST = SOSAT C – 88; SB = Soybean; CC = Commercial Control; N = Norman; C = Cettia CTA; AA = Atilagan Atilla.

significant at $p > 0.05$. The low values of moisture content in this study would enhance the storability and keeping quality of the products.

A proportionate increase in protein content occurred ranging between 6.97-1.56. The protein increase is due to supplementation with soybean which is known for its excellent source of protein because it contains all the essential amino acids, is very rich in minerals and is a good source of fat soluble vitamins (Alabi *et al.*, 2007; and Serrem

et al., 2011). Other studies have also reported a similar increase of protein content in soy-composite flours according to Singh *et al.* (2000) and Mashayekh *et al.* (2008). Sample C3 (80% whole wheat flour, 20% pearl millet, 0%) having the least which might be as a result of high percentage of pearl millet supplemented even though many researchers have proved its protein level to be comparable to other cereal grain. The highest value was observed in sample N4. All the samples were significantly different at

Table 4: Proximate Composition Funkaso Produced from Several Formulations

Formulations				Proximate Composition (%)							Energy (Kcal/100 g)
WT	ST	SB		Dry Matter	Moisture	Crude Protein	Crude Fat	Crude Fibre	Ash	CHO	
100CC	0	0	(CC)	86.30±0.00 ^{bcd}	13.69±0.00 ^{figh}	13.58±0.25 ^{abcde}	14.17±0.15 ^{hi}	2.86±0.02 ^{fg}	2.71±0.05 ^j	52.97±0.41 ^{defg}	393.78±0.62 ^{efg}
100AA	0	0	(A1)	86.45±0.23 ^{cdefg}	13.54±0.23 ^{defg}	10.86±3.21 ^{bcdefg}	14.43±0.50 ^j	2.88±0.01 ^g	2.79±0.01 ^k	51.47±3.48 ^{bcde}	395.29±3.46 ^{fg}
70AA	0	30	(A2)	87.09±0.21 ⁱ	12.90±0.02 ^b	18.32±1.21 ^{efg}	14.56±0.23 ⁱ	3.78±0.01 ^q	2.99±0.00 ^m	48.42±1.12 ^{abcd}	394.05±1.22 ^{efg}
80AA	20	0	(A3)	87.67±0.00 ^j	12.32±0.00 ^a	8.43±3.39 ^{defg}	15.02±0.00 ^j	2.51±0.01 ^d	2.96±0.00 ^l	50.74±3.40 ^{abcde}	403.92±0.06 ⁱ
56AA	14	30	(A4)	86.53±0.23 ^{cdefg}	13.40±0.20 ^{cdefg}	16.04±2.73 ^{defg}	12.49±0.00 ^d	3.60±0.01 ^{op}	3.87±0.00 ^q	50.59±2.89 ^{abcde}	378.97±0.83 ^a
70AA	30	0	(A5)	86.63±0.17 ^{cdefgh}	13.33±0.14 ^{bcd}	12.92±1.67 ^{abcde}	11.99±0.00 ^c	2.31±0.01 ^b	3.04±0.01 ^m	56.40±1.52 ^{ghi}	385.26±0.68 ^{cd}
49AA	21	30	(A6)	85.78±0.01 ^a	14.21±0.01 ⁱ	16.32±1.12 ^{abcde}	15.29±0.03 ^j	3.32±0.00 ^k	2.45±0.01 ^g	50.39±1.14 ^{abcde}	396.52±0.09 ^g
60AA	40	0	(A7)	86.89±0.00 ^{ghi}	13.10±0.00 ^{bcd}	10.83±2.11 ^{abcde}	16.31±0.01 ^{kl}	2.18±0.00 ^a	2.50±0.01 ^h	52.06±2.12 ^{def}	410.42±0.14 ^j
42AA	28	30	(A8)	86.40±0.00 ^{cde}	13.59±0.00 ^{efgh}	17.80±1.47 ^{defg}	13.00±0.00 ^c	3.21±0.00 ^j	3.46±0.01 ^o	50.92±1.49 ^{abcde}	383.92±0.12 ^{cd}
100C	0	0	(C1)	86.79±0.43 ^{efghi}	13.20±0.04 ^{bcd}	10.39±0.50 ^{ab}	14.18±0.15 ^{hi}	3.11±0.00 ⁱ	2.33±0.00 ^f	56.78±0.43 ^{ghi}	396.30±0.70 ^g
70C	0	30	(C2)	86.86±0.06 ^{efghi}	13.13±0.06 ^{bcd}	18.10±0.60 ^{defg}	16.99±0.00 ^j	3.54±0.02 ^{no}	2.59±0.01 ⁱ	47.63±0.55 ^{abc}	407.88±0.32 ^j
80C	20	0	(C3)	86.44±0.00 ^{cdefg}	13.55±0.00 ^{defg}	9.58±0.36 ^a	14.33±0.00 ^{hi}	3.02±0.00 ^h	2.22±0.00 ^c	57.28±0.35 ^{ghi}	396.45±0.04 ^g
56C	14	30	(C4)	86.69±0.00 ^{defghi}	13.20±0.10 ^{bcd}	17.64±0.78 ^{efg}	16.33±0.00 ^k	3.48±0.01 ^m	3.33±0.00 ⁿ	47.00±0.72 ^{ab}	401.58±0.37 ^{hi}
70C	30	0	(C5)	86.55±0.01 ^{cdefg}	13.44±0.01 ^{cdefg}	10.06±0.51 ^a	13.36±0.03 ^{ef}	2.67±0.00 ^c	2.01±0.01 ^c	58.43±0.45 ^{hi}	394.25±0.04 ^{efg}
49C	21	30	(C6)	86.43±0.00 ^{cdef}	13.63±0.02 ^{efgh}	17.50±0.35 ^{cdefg}	16.90±0.01 ^l	3.54±0.00 ^{no}	3.87±0.00 ^q	46.53±0.40 ^a	400.30±0.06 ^h
60C	40	0	(C7)	86.74±0.15 ^{defghi}	13.28±0.18 ^{bcd}	11.05±0.44 ^{abc}	10.96±0.04 ^b	2.41±0.00 ^c	2.12±0.00 ^d	60.16±0.62 ⁱ	383.50±0.67 ^{bc}
42C	28	30	(C8)	86.51±0.28 ^{cdefg}	13.52±0.29 ^{defg}	17.74±0.70 ^{efg}	16.49±0.16 ^k	3.40±0.03 ^l	3.04±0.02 ^m	46.79±0.40 ^{ab}	402.61±1.46 ^{hi}
100N	0	0	(N1)	86.42±0.12 ^{cdef}	13.64±0.02 ^{efgh}	13.59±0.59 ^{abcde}	10.69±0.03 ^{ab}	2.80±0.01 ^f	1.67±0.00 ^b	57.58±0.62 ^{hi}	380.96±0.15 ^{ab}
70N	0	30	(N2)	86.54±0.24 ^{cdefg}	13.46±0.24 ^{defg}	18.35±0.70 ^{fg}	12.37±0.01 ^d	3.61±0.01 ^p	2.68±0.00 ^j	49.52±0.63 ^{abcde}	382.82±0.94 ^{bc}
80N	20	0	(N3)	86.53±0.05 ^{cdefg}	13.46±0.05 ^{defg}	11.76±0.69 ^{abcd}	12.62±0.03 ^d	2.67±0.00 ^e	1.33±0.00 ^a	58.13±0.77 ^{hi}	393.22±0.16 ^{ef}
56N	14	30	(N4)	86.61±0.34 ^{cdefgh}	13.38±0.34 ^{cdefg}	18.31±0.07 ^{fg}	13.01±0.01 ^c	3.53±0.01 ^{mn}	3.34±0.01 ⁿ	48.41±0.37 ^{abcd}	384.01±1.33 ^{cd}
70N	30	0	(N5)	86.42±0.02 ^{cdef}	13.64±0.01 ^{efgh}	14.17±1.71 ^{abcde}	13.62±0.03 ^{fg}	2.49±0.05 ^d	2.01±0.01 ^c	54.04±1.73 ^{ef}	395.50±0.39 ^{fg}
49N	21	30	(N6)	86.99±0.01 ^{hi}	13.00±0.01 ^{bc}	18.48±0.01 ^{abcde}	13.68±0.00 ^{fg}	3.54±0.03 ^{no}	2.68±0.01 ^j	48.60±0.04 ^{abcd}	391.49±0.27 ^e
60N	40	0	(N7)	85.98±0.01 ^{ab}	14.01±0.01 ^{hi}	13.31±0.90 ^{abcde}	10.36±0.03 ^a	2.39±0.03 ^c	1.66±0.00 ^b	58.25±0.90 ^{hi}	379.56±0.35 ^a
42N	28	30	(N8)	86.20±0.01 ^{bc}	13.80±0.01 ^{gh}	17.04±0.01 ^{defg}	13.96±0.04 ^{gh}	3.39±0.03 ^l	3.62±0.04 ^p	49.18±0.12 ^{abcd}	386.53±0.05 ^d

Note: Means (±SE) in the same column having different superscripts are significantly ($p < 0.05$) different. WT = Wheat; ST = SOSAT C – 88; SB = Soybean; CC = Commercial Control; N = Norman; C = Cettia CTA; AA = Atillagan Atilla

$p > 0.05$ with the exception (C6, N6); (C2, N7); (A3, C5); (A4, C1) and (A1 and A7) respectively.

A value range of 1.88-8.67 fat content was obtained from these studies with the least recorded in sample N7 and the highest in sample C4. No significant difference in fat content were observed at $p < 0.05$ for these sample formulation A3, A5, A7; C1, C3, N3, N5; C8, A8; and C8, N2 respectively.

A high amount of crude fibre was observed in this study when compared to the amount in conventional wheat grains (local), these could be as a result of wholeness of the grain (wheat), varietal difference as the samples used in these studies were improved and might have affected the overall grain composition. The crude fibre content ranged between 2.42-4.22, significant variation was observed among the samples. The high crude fibre is most likely from the bran of the whole-wheat flour and the hull of soy beans, which represents variable fraction of dietary fibre and includes mostly the lignin, cellulose and hemicelluloses components (Mannay and Shadaksharaswamy, 2005). The least value was observed in sample A7 and highest in sample A2. Sample having the same formulation in variety C and N are not different at 5% level of significance.

An increasing trend was observed in the ash content level with pearl millet and soybean incorporation. The value ranged between 1.64-2.32%, least being observed in sample N1 (100% whole wheat) and the highest in N8 (42% whole wheat, 28% pearl millet, 30% soybean).

Value range of carbohydrate (60.59 to 75.55) was observed among all the flour samples. N7 had the highest value of 75.55 and the least sample was observed in sample N4 which was also found to be insignificantly different to C8 (60.70) at $p < 0.05$. Sample A (100%), A5 (70% whole wheat flour, 30% pearl millet, 0% soybean) and A3 (80% whole wheat flour, 20% pearl millet, 0%) an were also insignificantly different. A concomitant decrease in carbohydrate and increase in protein occurred, these could be as result of soybean flour inclusion as they are rich sources of protein. These trends have been reported by many researchers. The resulting total carbohydrate content in this study indicates that these types of flour are classified as food of the group one or food energy supplier of nutritive and economical value which could represent good sources for industrial flour and starch (FAO, 1998).

Functional Properties of *Tunkaso* Produced from Several Formulations

The result of the functional properties of *Funkaso* produced from different formulation is presented in Table 5, where water absorption ranged from 2.00-3.03 ml/g. The highest and lowest values were observed in C1 (100% Whole wheat flour) and N6 (49% whole wheat flour, 21% pearl millet, 30% soybean). Most samples that were not supplemented with soybean were insignificantly different ($p < 0.05$) except few. Considering the water absorption in terms of varietal difference, the CC (Control Commercial) was insignificantly different to formulation in variety A and C not containing soybean (A1, A3, A5, C1, C2 and N1). Sample C1 having higher water absorption when compared to the flour from which it was processed, and this might be as result of protein concentration, loss of conformal structure and leaching of amylose during processing (Butt and Batool, 2010). Sample N3, N4 and N5 were indifferent ($P < 0.05$).

A value range of 0.50-1.00 was observed in oil absorption of *funkaso*. Insignificant difference ($p < 0.05$) was observed in the control and most samples that are not supplemented with soybean in variety A and C, but variety N appeared to be different. Increased oil absorption was observed with incorporation of soybean in all the samples, similar result were reported by Gangale and Jadhao (2016). These could possibly be due to protein concentration and their conformational properties in foods which also influence oil absorption (Ige *et al.*, 1984; and Ahmad and Prakash, 2006).

The bulk density of *funkaso* flour increased concomitantly with supplementation at varying levels. The resulting value ranged between 0.71-1 ml/g. Bulk density give reflection/indication of relative volume of packaging material, mixing quality of particulate matter and load the flour sample accommodates if allowed to rest directly on the matter. The density of processed products dictate the characteristics of its container or package product, density influences the amount and strength of packaging material, texture or mouth feel (Apotiola *et al.*, 2016). According to Basman *et al.* (2003) higher bulk density is desirable for greater ease of dispensability of flours. In contrast, however, low bulk density would be an advantage in the formulation of complementary foods (Ugwu and Ukpabi, 2002).

Water swelling capacity of the *funkaso* ranged from 16-28 ml/10 g at 30 °C and 18-29.33 ml/10 g at 70 °C. Water swelling capacity decreased significantly at $p < 0.05$ for both

temperatures but lower in samples with soybean, and significantly different ($p < 0.05$) when compared to CC with other samples. High swelling capacity has been reported as part of the criteria for a good quality product (Apotiola

et al., 2016). Among all the samples at varying temperature sample A1 had the highest at 60 °C which is as result of increased temperature that enabled the starch to immobile water and swells (Ikegwu et al., 2009).

Table 5: Functional Properties of Funkaso from Different Formulation

Formulations (%)				Water Abs	Oil Abs	WSC 30 °C	WSC 70 °C	BD
WT	ST	SB		(ml/g)	(ml/g)	(g/ml)	(g/ml)	(g/ml)
100CC	0	0	(CC)	2.93±0.03 ^c	0.50±0.00 ^a	27.00±0.00 ^k	28.66±0.33 ^{lm}	0.89±0.00 ^k
100AA	0	0	(A1)	2.96±0.03 ^c	0.50±0.00 ^a	28.00±0.00 ^l	29.33±0.33 ⁿ	0.76±0.00 ^h
70AA	0	30	(A2)	2.53±0.03 ^{cd}	1.00±0.00 ^d	20.33±0.33 ^d	26.00±0.00 ^j	0.83±0.00 ^j
80AA	20	0	(A3)	3.00±0.00 ^e	0.50±0.00 ^a	22.33±0.33 ^h	20.33±0.33 ^c	0.83±0.00 ^j
56AA	14	30	(A4)	2.50±0.11 ^{cd}	1.00±0.00 ^d	21.33±0.33 ^{ef}	22.33±0.33 ^{fg}	0.76±0.00 ^h
70AA	30	0	(A5)	3.00±0.00 ^e	0.50±0.00 ^a	21.66±0.33 ^{fg}	24.00±0.00 ^h	0.71±0.00 ^d
49AA	21	30	(A6)	2.53±0.03 ^{cd}	0.50±0.00 ^a	19.33±0.33 ^c	21.00±0.00 ^{cd}	0.71±0.00 ^d
60AA	40	0	(A7)	2.33±0.33 ^{bc}	1.00±0.00 ^d	21.00±0.00 ^e	21.33±0.33 ^{de}	0.66±0.00 ^b
42AA	28	30	(A8)	2.50±0.00 ^{cd}	0.50±0.05 ^a	20.33±0.33 ^d	22.66±0.33 ^{fg}	1.00±0.00 ⁿ
100C	0	0	(C1)	3.03±0.03 ^e	0.73±0.03 ^b	24.33±0.33 ^j	28.33±0.33 ^l	0.76±0.00 ^h
70C	0	30	(C2)	2.96±0.03 ^e	0.96±0.03 ^d	16.00±0.00 ^a	21.00±0.00 ^{cd}	0.76±0.00 ^h
80C	20	0	(C3)	2.20±0.00 ^{ab}	0.50±0.00 ^a	20.00±0.00 ^d	19.00±0.00 ^b	0.82±0.00 ⁱ
56C	14	30	(C4)	2.23±0.03 ^{ab}	1.00±0.00 ^d	19.33±0.33 ^c	18.33±0.33 ^{ab}	0.71±0.00 ^e
70C	30	0	(C5)	2.50±0.00 ^{cd}	0.50±0.00 ^a	23.00±0.00 ⁱ	22.33±0.33 ^{fg}	0.71±0.00 ^d
49C	21	30	(C6)	2.23±0.03 ^{ab}	0.93±0.03 ^{cd}	16.33±0.33 ^a	18.00±0.00 ^a	0.76±0.00 ^g
60C	40	0	(C7)	2.70±0.11 ^d	0.50±0.00 ^a	22.00±0.00 ^{gh}	24.33±0.33 ^{hi}	0.71±0.00 ^c
42C	28	30	(C8)	2.53±0.03 ^{cd}	1.00±0.00 ^d	20.00±0.00 ^d	22.00±0.00 ^{ef}	0.99±0.00 ^m
100N	0	0	(N1)	3.00±0.00 ^e	1.00±0.00 ^d	23.00±0.00 ⁱ	27.00±0.00 ^k	0.92±0.00 ^l
70N	0	30	(N2)	2.50±0.00 ^{cd}	0.73±0.03 ^b	23.00±0.00 ⁱ	24.66±0.33 ^{hi}	0.76±0.00 ^h
80N	20	0	(N3)	2.30±0.00 ^{bc}	0.50±0.00 ^a	22.00±0.00 ^{gh}	26.00±0.00 ^j	0.71±0.00 ^c
56N	14	30	(N4)	2.43±0.03 ^{bc}	1.00±0.00 ^d	21.00±0.00 ^e	24.00±0.00 ^h	0.71±0.00 ^d
70N	30	0	(N5)	2.30±0.00 ^{bc}	0.90±0.00 ^c	19.00±0.00 ^c	25.00±0.00 ⁱ	0.62±0.00 ^a
49N	21	30	(N6)	2.00±0.00 ^a	1.00±0.00 ^d	18.00±0.00 ^b	22.33±0.33 ^{fg}	0.76±0.00 ^h
60N	40	0	(N7)	2.53±0.03 ^{cd}	1.00±0.00 ^d	18.00±0.00 ^b	22.66±0.33 ^{fg}	0.71±0.00 ^d
42N	28	30	(N8)	2.20±0.00 ^{ab}	0.76±0.06 ^b	19.00±0.00 ^c	23.00±0.00 ^g	0.72±0.00 ^f

Note: Means (±SE) in the same column having different superscripts are significantly ($p < 0.05$) different. WT = Wheat; ST = SOSAT C – 88; SB = Soybean; CC = Commercial Control; N = Norman; C = Cettia CTA; AA = Atillagan Atilla.

Sensory Scores of Funkaso Produced from Various Formulations

The Sensory characteristics of the *funkaso* presented in Table 6 shows the overall acceptability of prepared *funkaso*

was defined by sensory evaluation which was performed by 15 number of semi- trained panelist. The result from these study shows that, there is significant difference ($p>0.05\%$) in overall acceptability, appearance, taste,

Table 6: Sensory Scores of Funkaso Produced from Various Formulations¹

Formulations				Sensory Panel Scores ²				
WT	ST	SB		Appearance	Taste	Texture	Colour	Overall
100CC	0	0	(CC)	7.83±0.16 ^g	6.08±0.67 ^{abcde}	6.08±0.60 ^{abcde}	6.83±0.69 ^{def}	7.33±0.18 ^{hi}
100AA	0	0	(A1)	7.83±0.20 ^g	8.33±0.25 ^f	7.75±0.32 ^h	8.16±0.27 ^f	8.16±0.16 ⁱ
70AA	0	30	(A2)	6.41±0.54 ^{bcdefg}	6.41±0.63 ^{bcde}	7.00±0.40 ^{efgh}	7.25±0.27 ^{ef}	6.75±0.60 ^{defgh}
80AA	20	0	(A3)	6.25±0.56 ^{abcdef}	5.58±0.46 ^{abcde}	6.58±0.37 ^{bcdefgh}	6.66±0.33 ^{cde}	6.16±0.54 ^{abcde}
56AA	14	30	(A4)	6.58±0.43 ^{bcdefg}	5.75±0.49 ^{abcde}	6.91±0.43 ^{defgh}	6.91±0.37 ^{def}	5.75±0.37 ^{abcde}
70AA	30	0	(A5)	6.75±0.30 ^{cdefg}	5.50±0.52 ^{abcde}	5.50±0.50 ^{abcd}	6.33±0.37 ^{abcde}	6.08±0.45 ^{abcde}
49AA	21	30	(A6)	6.83±0.36 ^{cdefg}	5.58±0.43 ^{abcde}	6.25±0.44 ^{abcde}	6.91±0.39 ^{def}	6.66±0.41 ^{bcde}
60AA	40	0	(A7)	6.58±0.31 ^{bcdefg}	6.91±0.37 ^e	7.25±0.32 ^{gh}	6.66±0.37 ^{cde}	6.91±0.35 ^{efghi}
42AA	28	30	(A8)	7.41±0.31 ^{fg}	6.50±0.19 ^{cde}	6.75±0.30 ^{cdefgh}	6.75±0.30 ^{de}	7.08±0.31 ^{fghi}
100C	0	0	(C1)	6.91±0.52 ^{defg}	6.75±0.59 ^{de}	7.16±0.44 ^{fgh}	7.08±0.28 ^{def}	7.16±0.44 ^{ghi}
70C	0	30	(C2)	6.83±0.40 ^{cdefg}	5.91±0.33 ^{abcde}	6.33±0.51 ^{abcde}	6.66±0.22 ^{cde}	6.58±0.22 ^{bcde}
80C	20	0	(C3)	6.66±0.28 ^{bcdefg}	6.25±0.49 ^{bcde}	6.08±0.43 ^{abcde}	6.50±0.35 ^{bcde}	6.50±0.41 ^{bcde}
56C	14	30	(C4)	6.50±0.37 ^{bcdefg}	5.33±0.39 ^{abcde}	6.00±0.32 ^{abcde}	5.83±0.34 ^{abcde}	5.66±0.33 ^{abcde}
70C	30	0	(C5)	6.58±0.43 ^{bcdefg}	6.33±0.35 ^{bcde}	5.08±0.28 ^{ab}	6.00±0.34 ^{abcde}	5.58±0.14 ^{abcde}
49C	21	30	(C6)	6.33±0.37 ^{abcde}	5.66±0.33 ^{abcde}	5.66±0.51 ^{abcde}	6.83±0.20 ^{def}	6.16±0.40 ^{abcde}
60C	40	0	(C7)	5.91±0.48 ^{abcde}	5.83±0.40 ^{abcde}	5.58±0.51 ^{abcde}	6.00±0.46 ^{abcde}	5.83±0.40 ^{abcde}
42C	28	30	(C8)	5.41±0.52 ^{abc}	5.33±0.33 ^{abcde}	5.41±0.43 ^{abc}	5.83±0.51 ^{abcde}	5.66±0.51 ^{abcde}
100N	0	0	(N1)	5.41±0.39 ^{abc}	4.50±0.48 ^a	5.75±0.46 ^{abcde}	4.91±0.41 ^a	5.41±0.49 ^{abcde}
70N	0	30	(N2)	5.25±0.44 ^{ab}	5.16±0.61 ^{abcd}	5.00±0.47 ^a	5.08±0.49 ^{ab}	5.16±0.53 ^{abc}
80N	20	0	(N3)	5.00±0.50 ^a	5.00±0.53 ^{abc}	5.08±0.64 ^{ab}	5.16±0.56 ^{ab}	5.00±0.57 ^{ab}
56N	14	30	(N4)	5.66±0.43 ^{abcd}	5.08±0.60 ^{abcd}	5.33±0.39 ^{abc}	5.91±0.46 ^{abcde}	5.25±0.55 ^{abcd}
70N	30	0	(N5)	5.83±0.40 ^{abcde}	5.25±0.50 ^{abcde}	5.33±0.35 ^{abc}	5.25±0.42 ^{abc}	4.83±0.38 ^a
49N	21	30	(N6)	6.58±0.43 ^{bcdefg}	4.75±0.60 ^{ab}	5.50±0.33 ^{abcd}	5.91±0.51 ^{abcde}	5.33±0.48 ^{abcd}
60N	40	0	(N7)	5.91±0.66 ^{abcde}	5.66±0.49 ^{abcde}	5.25±0.42 ^{abc}	5.66±0.41 ^{abcd}	5.50±0.46 ^{abcde}
42N	28	30	(N8)	7.16±0.29 ^{efg}	6.58±0.64 ^{cde}	5.91±0.35 ^{abcde}	6.16±0.63 ^{abcde}	5.91±0.67 ^{abcde}

Note: ¹ Means (±SE) in the same column with different superscripts are significantly ($p< 0.05$) different. ² Scoring system using the 9-point hedonic scale in which 9-like extremely and 1-dislike extremely. WT = Wheat; ST = SOSAT C – 88; SB = Soybean; CC = Commercial Control; N = Norman; C = Cettia CTA; AA = Atillagan Atilla.

colour and texture evaluated. The highest score in terms of taste, texture, colour and overall acceptability were observed in sample A1 with 8.33, 7.75, 8.16 and 8.16 respectively. The score for overall acceptability ranged between 4.83 and 8.16, A1 is significantly different ($p > 0.05$) to all the samples, and no significant difference were observed in sample A3, A5, C6 and N8; and A7, N1 and N5 among others but differ with the CC. Sample N5 had the least score. A score range of 4.75 to 8.33, 4.91 to 8.16, 5.00 and 7.75 was observed in terms of taste, colour and texture, A1 is significantly different from CC at $P < 0.05$ while all other samples were not significantly different. No significant difference was observed in terms of Appearance with a range from 5.00 and 7.83, with the exception of A1 and CC. Sample A1 remains the most acceptable following the various parameters used in evaluating its acceptability. The score reduces with the supplementation of pearl millet and soybeans, and this is in line with the finding of Apotiola (2013) where cookies was supplemented with yam and soybean.

In terms taste and texture samples CC, A3, A4, A5, A6, C2, C4, C6, C7, C8, N5 and N7; and CC, A6, C3, C4 and N8 are insignificantly different at 5% level, also no significant difference was observed in colour CC, C1, A4 and A6; and A5, C4, C5, N4, N6 and N8 at $p < 0.05$ respectively.

CONCLUSION

Funkaso flour blends were produced from mixtures of wheat cultivars, pearl millet, soybean along with other ingredients such as baking powder, yeast, salt and water. *Funkaso* produced from these blends were subjected to proximate composition, functional properties and sensory evaluation. Protein and fat contents increased significantly ($P < 0.05$) with soybean supplementations. Addition of soybean increased the protein and fat contents of complementary food significantly ($P < 0.05$). Sensory scores tasted by panelists were generally high and therefore wheat can be supplemented with soybean and pearl millet at 30% and 28% levels, respectively in *funkaso* processing which had increased the protein and fat contents which could also increase the profit margin and save foreign exchange earnings by reducing importation of wheat.

ACKNOWLEDGMENT

Mrs. Fannah Mustapha Adam is grateful to the Management of University of Maiduguri for their financial support.

REFERENCES

- Adejuyitan JA, Otunola ET, Akande EA, Bolarinwa IF and Oladokun F M (2009), "Some Physicochemical Properties of Flour Obtained from Fermentation of Tigernut (*Cyperus esculentus*) Sourced from a Market in Ogbomoso, Nigeria", *African Journal of Food Science*, Vol. 3, No. 2, pp. 51-55.
- Ahmadzadeh G and Prakash J (2006), "Effects of Germination and Dehulling on Functional Properties of Legume Flours", *Journal of Science of Food and Agriculture*, Vol. 86, No. 8, pp. 1189-1195, doi: 10.1002/jsfa.2460.
- Alabi M O and Anuonye J C (2007), "Nutritional Sensory Attributes of Soy Supplemented Cereal Meals", *Nigeria Food Journal*, Vol. 25, No. 1, pp. 100-110.
- Aminu K M, Ajayi D, Ikwelle M C and Anaso AB (1998), "Trends in Millet Production in Nigeria", pp. 41-49.
- Annapure U S, Singhal R S and Kulkarni P R (1999), "Screening of Hydrocolloids for Reduction in Oil Uptake of a Model Deep-Fat Fried Product", *FETT/Lipid*, Vol. 101, pp. 217-221.
- Anyinka J U and Uwaegbute A C (2005), "Frequency of Consumption and Nutrient Content of Some Snacks Eaten by Adolescent Female Secondary and University Students in Abia State", *Nigerian Journal Nutritional Science*, Vol. 26, No. 2, pp. 10-15.
- AOAC (1990), "Association of Official Analytical Chemist", *Official Method of Analysis*, 14th Edition, Washington DC.
- AOAC (2006), "Association of Official Analytical Chemists", *Official Method of Analysis*, W Horwitz (Ed.), 18th Edition, Washington DC.
- AOAC (2013), *Official Method of Analysis of Association of Analytical Chemist*, 16th Edition, pp. 6-10, Washington DC.
- Apotiola Z O and Fashakin J F (2013), "Evaluation of Cookies from Wheat Flour, Soybean Flour and Cocoyam Flour Blends", *Food Science and Quality Management*, Vol. 14.
- Badau M H and Magaji AM (2011), "Mineral Content and Acceptability of Wheat Alkaki with Added Pearl Millet and Cowpea Flour", Proceedings of 35th Annual Conference AGM Organized by NIFST, held on 10-14, October, at Benue Hotels Ltd., Makurdi.

- Badau MH, Jidaeni IA and Nkama I (2005), "Production, Acceptability and Microbiology Evaluation of Weaning Food Formulations", *Journal of Tropical Pediatrics*, Vol. 52, No. 3, pp. 166-172.
- Badau MH, Bimba S and Kyari F Y (2009), "Quality of Rice Garabian as Affected by Addition of Pearl Millet Flour", Proceedings of 33rd Annual Conference AGM of NIFST, Held on 12-16 October at Banquet Hall Government House Yola, Adamawa State, Nigeria.
- Badau MH, Ngozi C and Danbaba N (2013), "Quality of Garabia (ANigeria Traditional Snack) from Four Varieties of Rice as Affected by the Additional of Cowpea", *Advanced Journal of Food Science and Technology*, Vol. 5, No. 3, pp. 249-254.
- Badau MH, Silo S J and Usman Z L (2008), "Quality of Wheat Alkaki as Affected by Addition of Pearl Millet Flour", Proceedings of 32nd Annual Conference AGM of NIFST, Held on 13-17 October, at Senate Building, Ladoke Akintola University of Technology, Ogbomosho, Nigeria.
- Basman A, Koksel H and Ng P K W (2003), "Utilization of Transglucanase Use to Increase the Level of Barley and Soyflour Incorporation in Wheat Flour Breads", *J. Food Sci.*, Vol. 68, No. 8, pp. 2453-2460.
- Belderok Robert 'Bob', Mesdag Hans and Donner Dingena A (2000), "Bread-Making Quality of Wheat", Springer, p. 3, ISBN: 0-7923-6383-3.
- Benitez L V (1989), "Amino Acid and Fatty Acid Profiles in Aquaculture Nutrition Studies", in S S De Silva (Ed.), *Fish Nutrition Research in Asia A Proceedings of the Third Asian Fish Nutrition Network Meeting*, Asian Fish Society Special Publication, Asian Fisheries Society, Manila, Philippines.
- Beuchat L R (1977), "Functional and Electrophoretic Characteristics of Succinylated Peanut Flour Protein", *J. Agric. Food Chem.*, Vol. 25, pp. 258-261.
- Bicar E H, Woodman-Clikeman W, Santong V, Peterson J M, Yang S S, Lee M and Scott M P (2008), "Transgenic Maize Endosperm Containing a Milk Protein has Improved Amino Acid Balance", *Transgenic Research*, Vol. 17, pp. 59-71.
- Bidinger F R and Hash C T (2003), "Pearl Millet in 'Integration of Physiology and Molecular Biology in Plant Breeding'", H Nguyen and A Blum (Eds.), Marcel Dekker (in Press), New York.
- Blanco J L and Sherbo G H (1978), *Standard Methods for the Examination of Dairy Products*, American Public Health Association, N.W.D.C.
- Butt M S and Batool R (2010), "Nutritional and Functional Properties of Some Promising Legumes Protein Isolates", *Pakistan Journal of Nutritional*, Vol. 9, No. 4, pp. 373-379.
- CGIAR (2011), "The Revival of Traditional Dry-Land Crops", www.cgiar.org/news/the-revival-of-traditional-dryland-crops
- Chibuzo E C and Ali H (1994-95), "Sensory and Physicochemical Evaluation of Millet-Groundnut Blend as Weaning Food", *Annals of Borno*, Vol. 11, No. 12, pp. 181-190.
- Chopra V L (2001), "Breeding Field Crops", IBH Publishing Co Pvt Ltd., Oxford.
- Cornell H (2003), in: Cauvain S P (Ed.), "Bread Making: Improving Quality", Wood Head Publishing, Cambridge.
- Curtis, Rajaraman and Mac Pherson (2002), "Bread Wheat", Food and Agriculture Organization of the United Nations.
- Dahiya S and Kapoor A C (1994), "In Vitro and in Vivo Availability of Iron from Home Processed Supplementary Foods", *Journal of Food Science and Technology*, Vol. 31, No. 2, pp. 122- 125.
- Dhingra S and Jood S (2002), "Physico-Chemical and Nutritional Properties of Cereal-Pulse Blends for Bread Making", *Nutr. Health*, Vol. 16, No. 3, pp. 183-194.
- Doell B H, Ebdon C J and Smith C A (1981), "Trypsin Inhibitor Activity of Conventional Which are Part of the British Diet and Some Soy Products", *Plant Foods for Human Nutrition*, Vol. 13, pp. 139-150.
- Duncan D E (1955), "Multiple Range and Multiple F-tests", *Biometrics*, Vol. 11, pp. 1-42.
- Ejeta G, Hansen M M and Mertz E T (1987), "In Vitro Digestibility and Amino Acid Composition of Pearl Millet (*Pennisetum typhoides*) and Other Cereals", Proceedings of National Academy of Sciences (USA), Vol. 84, pp. 6016-6019.
- Elleuch M, Bedigian D, Roiseux O, Besbes S, Blecker C and Attia H (2011), "Dietary Fibre and Fibre-Rich by-Products of Food Processing: Characterisation,

- Technological Functionality and Commercial Applications”, *Rev. Food Chem.*, Vol. 124, pp. 411-421.
- El-Shenawy M, El-Shenawy M, JordiManes and Jose M S (2011), “*Listeria spp.* in Street-Vended Ready-to-Eat Foods”, *Interdisciplinary Perspectives on Infectious Diseases*, Vol. 28, No. 15, pp. 24-29.
 - EM M A(1973), “Vegetable Oils and Fats, their Production and Commercial Extraction”, Information Division, Unilever Ltd., England.
 - Falola AO, Olatidoye O P, Balogun I O and Opeifa A O (2011), “Microbiological Quality Analysis of Meat Pies Sold by Street Hawkers: A Case Study of Mainland Local Government Area of Lagos, Nigeria”, *Journal of Medical and Applied Biosciences*, Vol. 2, pp. 1-8.
 - FAO (1998), “Carbohydrates in Human Nutrition (Food and Nutrition Rome)”, p. 66.
 - FAO (2002), *World Agriculture: Towards 2015/2030*, Summary Report, Rome.
 - FAO (2007), Food and Agricultural Organization, Annual Publication, Rome, Italy.
 - Food and Agricultural Organization (2005), “Statistical Databases”, available at <http://apps.fao.org.FAOSTat>., Retrieved 27 January.
 - Food and Agricultural Organization (2015), “Statistical Databases”, available at <http://apps.fao.org.FAOSTat>., Retrieved 27 January.
 - Food Slavin J L, Jacobs D and Marquart L (2000), “Grain Processing and Nutrition”, *Crit Rev Food Sci Nutr.*, Vol. 40, pp. 309-326.
 - Gangale R A and Jadhao V G (2016), “Effect of Hydrocolloid on Indian Traditional Food Puri Bended with Pearl Millet and Soyabean Flour”, *International Journal of Innovations in Engineering and Technology (IJJET)*, Vol. 7, pp. 269-274.
 - Gibney A, Butler F and Dwyer E (1999), “Rheology and Adhesion of Fish Batter Coating Made from Flour from Irish Grown Wheat Varieties”, *Irish Journal of Agricultural and Food Research*, Vol. 38, pp. 241-249.
 - Gomez K and Gomez AA (1983), *Statistical Procedures for Agricultural Research*, 2nd Edition, John Wiley and Sons, New York.
 - Han J, Lee M and Lim S (2007), “Utilization of Oxidized and Cross-Linked Corn Starches in Wheat Flour Batter”, *Cereal Chemistry*, Vol. 84, pp. 582-586.
 - Henry W A and Morrison F B (1923), “Feeds and Feeding: A Handbook for the Student and Stockman”, The Henry-Morrison Co. Madison, Wisconsin, USA.
 - Henshaw R B and Agunbiade M O (2004), “Food Oils and Fats Technology: Utilization and Nutrition”.
 - Hill G M and Hanna W W (1990), “Nutritive Characteristics of Pearl Millet Grain in Beef Cattle Diets: Issues in New Crops and New Uses”, *Journal of Animal Sciences*, Vol. 68, pp. 2061-2066.
 - Hosoney R C, Andrews D J and Clark H (1987), “Sorghum and Pearl Millet”, in: *Nutritional Quality of Cereal Grains: Genetic and Agronomic Improvement*, Vol. 28, pp. 397-456, ASA, Monograph.
 - Hussein K M Ashraf and Ashraf M Y (2008), “Relationship Between Growth and Ion Relation in Pearl Millet (*Pennisetum glaucum*) at Different Growth Stages Under Salt Stress”, *African Journal of Plant Science*, Vol. 2, No. 3, pp. 23-27.
 - ICRISAT/FAO (1996), “The World Sorghum and Millet Economies: Facts, Trends and Outlook”, *ICRISAT*, p. 68, Patancheru, India and FAO, Rome.
 - Iden N U A and Showemimo F A (2004), “Cereal Crops of Nigeria”, *Principles of Production and Utilization*, Vol. xxii, p. 337, Ade Commercial Printing Press, Ibadan, Nigeria.
 - Igbabul B D, Idikwu H O and Inyang C U (2012), “Effect of Fermentation on Some Functional Properties of *Mucunasloanei* and *Detariummicrocarpum*”, *J. of Food Technol.*, Vol. 10, No. 3, pp. 83-86.
 - Ige M M, Ogunsua A O and Oke O L (1984), “Functional Properties of the Proteins of Some Nigerian Oil Seeds: Conophor Seeds and Three Varieties of Melon Seeds”, *Agric and Food Chem.*, Vol. 31, pp. 822-825.
 - Ihekoronye A I and Ngoddy P O (1985), *Integrated Food Science and Technol. for the Tropics*, 1st Edition, pp. 261, 265, 291, McMillan Publishers.
 - Ihekoronye A I and Ngoddy P O (1985). “Integrated Food Science and Technology for Tropics”, pp. 24-251, Macmillanpub, London.

- IITA (1979), *Soil and Plant Analysis*, International Institute for Tropical Agriculture, Manual, Series No. 1, IITA, Ibadan.
- Ikegwu O J, Nwobasi V, Odoh M O and Oledinma N U (2009), "Evaluation of the Pasting and Some Functional Properties of Starch Isolated from Some Improved Cassava Varieties in Nigeria", *Afri. J. Biotechnology*, Vol. 8, No. 10, pp. 2310-2315.
- Ingbian E K and Akpapunam M A (2005), "Appraisal of Traditional Technologies in the Processing and Utilisation of Mumu, a Cereal Based Local Food Product", *African Journal of Food, Agriculture, Nutrition and Development*, Vol. 5, No. 2, pp. 11-15.
- IRD (2009), Institut de recherche pour le développement, Pearl Millet, Food for the Future in the Sahel, *Scientific Bulletin*, p. 325.
- Iwe M O (2003), "The Science and Technology of Soybeans Chemistry, Nutrition, Processing, Utilization", Rojoint Communications Services Ltd., Umuahia, Nigeria.
- Izge A U (2006), "Combining Ability and Heterosis of Grain Yield Components Among Pearl Millet (*Pennisetum glaucum* L. R. Br) in Breeds", Ph.D. Thesis, p. 148, Federal University of Technology, Yola, Nigeria.
- Jay J M (2000), *Modern Food Microbiology*, 6th Edition, p. 168, Aspen Publishers, Inc Gaithersburg, Maryland.
- Kamaljit K, Baljeet S and Amarjeet K (2013), "Preparation of Bakery Products and Incorporating Pea Flour as a Functional Ingredients Department of Food Science and Technology, Punjab Agricultural University, Ludhlana, India", *American Journal of Food Technology*, Vol. 5, pp. 130-135, 2010.
- Kent N L (1983), *Technology of Cereals*, 3rd Edition, Toronto Sydney, Paris Frankfurt, Pergam on Press, Oxford, New York.
- Khairawal I S, Rai K N, Andrew D J and Harnarayana A (1999), "Pearl Millet Breeding", Oxford and IBH Publishing Co. PVT Ltd.
- Ku K H, Park D J and Kim S H (1996), "Characteristics and Application of Soybean Hull Fractions Obtained by Microparticulated/Air-Classification", *Korean Journal of Food Science Technology*, Vol. 28, pp. 506-513.
- Kure O A, Bahago E J and Daniel E A (1998), "Studies on the Proximate Composition and Effect of Flour Particle Size of Acceptability of Biscuits Produced from Blends of Soybeans and Plantain Flours", *Namoda Tech. Scope J.*, Vol. 3, No. 2, pp. 17-22.
- Kwaw E and Sackey A S (2013), "Nutritional and Sensory Analysis of Millet Based Sponge Cake", *International Journal of Nutrition and Food Sciences*, Vol. 2, No. 6, pp. 287-293.
- Lee C C, Hoseney R C and Variona-Martson E (1982), "Development of Laboratory-Scale Single-Stage Cake Mix", *Cereal Chem.*, Vol. 59, pp. 389-392.
- Lee J S, Kim B K, Kim K H and Park D J (2008), "Preparation of Low-Fat Uptake Doughnut by Dry Particle Coating Technique", *Journal of Food Science*, Vol. 73, pp. 137-142.
- Liener I E (1989), "Legumes, Chemistry, Technology and Human Nutrition", Marced, Inc. Dekker, New York.
- Loewe R (1993), "Role of Ingredient in Batter Systems", *Cereal Foods World*, Vol. 38, pp. 673-677.
- Mannay S and Shadaksharaswamy C M (2005), *Foods: Facts and Principles*, 2nd Edition, New Age International Ltd. Publishers, New Delhi, India.
- Mashayekh M, Mahmoodi M R and Enterazzi M H (2008), "Effect of Fortification of Defatted Soy Flour on Sensory and Rheological Properties of Wheat Bread", *Int. J. Food Sci. Technol.*, Vol. 43, pp. 1693-1698.
- Mbah M, Ogban G I, Konlack G D, Useh M F and Asuquo A E (2012), "The Bacteriological Status of Five Selected Street Vended Cooked Foods in Calabar, Nigeria", *Journal of Pharmacy and Biological Sciences*, Vol. 2, No. 4, pp. 25-29.
- Mensah P, Nicholas S L and Mithani C L (2002), "Street Foods in Accra, Ghana: How Safe are They", *Bulletin of the World Health Organization*, Vol. 80, No. 7, pp. 546-554.
- Meyers M A (1990), "Functionality of Hydrocolloids in Batter Coating System", in: Kulp K (Ed.).
- Murphy P A (1982), "Phytoestrogen Content of Processed Soyabean Products", *Technology*, Vol. 36, pp. 60-64.
- Murty B R, Arunachalam V and Saxena M B L (1967), "Classification and Catalogue of a World Collection of

- Cultivated Sorghums and Pennisetum”, *Indian Journal of Genetics and Plant Breeding*, Vol. 27, pp. 1-74.
- National Research Council (1996), “Lost Crops of Africa”, Vol. 1, p. 583, Grains, National Academy Press, Washington DC.
 - Neill Richard (2002), “Booze: The Drinks Bible for the 21st Century”, p. 112, Octopus Publishing Group–Cassell Illustrated, ISBN: 1-84188-196-1.
 - Niba L L (2003), “Processing Effects on Susceptibility of Starch to Digestion in Some Dietary Starch Sources”, *International Journal of Food Science and Nutrition*, Vol. 54, pp.97-109.
 - Nkama I (1991), “Traditional Method of Production of High Protein Food from Grain Legumes in the North-Eastern Nigeria”, Paper Presented at the International Seminar on the Development of High Protein Energy Food from Grain Legumes Organised by the Association of African University (A.A.U), Ghana, 5-7 February.
 - Nwokolo E (1996), “Bambara Groundnuts (*Vigna subterranean*)”, in: *Food and Feed from Legumes and Oil Seeds*, Nwokolo E and Smart J (Eds.), pp. 216-221, Chapman and Hall, London.
 - Nyenje E M, Collins E, Odjadjare Noline F, Tanih Ezekiel G and Roland N N (2012), “Foodborne Pathogens Recovered from Ready-to-Eat Foods from Roadside Cafeterias and Retail Outlets in Alice, Eastern Cape Province, South Africa: Public Health Implications”, *International Journal of Environmental Research and Public Health*, Vol. 9, pp. 2608-2619.
 - Obilana A B, Manyasa E Millets (2002), in “*Pseudocereals and Less Common Cereals: Grain Properties and Utilization Potential*”, P S Belton and J R N Taylor (Eds.), Springer-Verlag, pp. 177-217, Berlin, Heidelberg, New York.
 - Odepidan V and Ilo M (1996), “An Outline of Microbiology Techniques: A Laboratory Manual”, pp. 8-83, Emman Food Science, and Technology Department Publications, Ibadan.
 - Ohiokpehai O (2003a), “Food Processing and Nutrition: A Vital Link in Agricultural Development”, *Pakistan Journal of Nutrition*, Vol. 2, No. 3, pp. 204-207.
 - Ojediran J O, Adamu M A and Jim-George D L (2010), “Some Physical Properties of Pearl Millet (*Pennisetum glaucum*) Seeds as a Function of Moisture Content”, *African Journal of General Agriculture*, Vol. 6, p. 1.
 - Ojinnaka M C, Ebinyasi C S, Ihemeye A and Okorie S U (2013), “Nutritional Evaluation of Complementing Food Gruel Formulated from Blends of Soybean Flour and Ginger Modified Cocoyam Starch”, *Advance Journal of Food Science and Technology*, Vol. 5, No. 1, pp. 1325-1330.
 - Okaka J C and Potter N N (1979), “Physicochemical and Functional Properties of Cowpea Powders Processed to Reduce Beany Flavor”, *J. Food Sci.*, Vol. 44, pp. 1235-1240.
 - Okoye J I, Ezigbo V O and Animalu I L (2010), “Development and Quality Evaluation of Weaning Foods Fortified with African Yam Bean Flour Cont”, *J. Agric Sci.*, Vol. 4, pp. 1-6.
 - Omidiran A T, Sanni L O, Sobukola O P, Adebowale A A, Obadina A O, Kajihaua O E, Adegunwa M O, Tomlins K and Wolfgang T (2015), “Evaluation of Some Properties of Wheat-Brewers’ Spent Cassava Flour Blends”, *Nigerian Food Journal*, Vol. 33, pp. 10-18.
 - Onwuka G I (2005), *Food Analysis and Instrumentation, Theory and Practice*, Naphthali Prints, A Division of HG Support Nig. Ltd., Lagos, Nigeria.
 - Palmer John J (2001), “How to Brew”, p. 233, Defenestrative Pub Co., ISBN: 0-9710579-0-7.
 - Penfield M P and Campbell A M (1990), *Experimental Food Science*, 3rd Edition, Academy Press, Inc. San Diego, New York.
 - Rai K N, Murty D S, Andrews D J and Bramel-Cox P J (1999), “Genetic Enhancement of Pearl Millet and Sorghum for the Semi-Arid Tropics of Asia and Africa”, *Genome*, Vol. 42, pp. 617-628.
 - Rooney L W and McDonough C M (1987), “Food Quality and Consumer Acceptance of Pearl Millet”, in Proceedings of the International Pearl Millet Workshop 7-11 April 1986, in J R Witcombe and S R Beckerman (Eds.), pp. 43-61, ICRISAT, Patancheru, India.
 - Salunkhe D K, Chavan J C, Adsule R N and Kadam S S (1992), “World Oil Seeds”, *Chemistry, Technology and Utilization on An ÀV Book Published*, Van Nostrand, New York.

- Serna-Saldivar S (2010), "Cereal Grains: Properties", Processing and Nutritional Attributes, p. 535, Retrieved 22 June 2015.
- Serrem C, Kock H and Taylor J (2011), "Nutritional Quality, Sensory Quality and Consumer Acceptability of Sorghum and Bread Wheat Biscuits Fortified with Defatted Soy Flour", *Int. J. Food Sci. Technol.*, Vol. 46, pp. 74-83.
- Shewry PR, Tathan AS and Kasarda DD (1992), "Cereal Proteins in Coeliac Disease", in *Coeliac Disease*, M N Marsh (Ed.), Blackwell Science, Oxford.
- Singh R, Singh G and Chauhan G S (2000), "Nutritional Evaluation of Soy Fortified Biscuits", *J. Food Sci. Technol.*, Vol. 37, pp. 162-164.
- Singh S, Ria C S and Saxena D C (2008), "Effect of Incorporating Sweet Potato Flour to Wheat Flour on the Quality Characteristics of Cookies", *Afr. J. Food Sci.*, Vol. 2, pp. 65-72, University Press, Cambridge, England.
- Slavin J L (2005), "Dietary Fiber and Body Weight", *Nutr.*, Vol. 21, pp. 411-418.
- Slavin J L, Jacobs D and Marquart L (2000), "Grain Processing and Nutrition", *Crit Rev Food Sci Nutr.*, Vol. 40, pp. 309-326.
- Ugwu B O and Ukpabi U J (2002), "Potential of Soy-Cassava Flour Processing to Sustain Increasing Cassava Production in Nigeria", *Outlook on Agriculture*, Vol. 31, No. 2, pp. 129-135.
- Wang X S, Tang C H, Li B S, Yang X Q, Li L and Ma C Y (2008), "Effect of High Pressure Treatment on Some Physico-Chemical Functional Properties of Soy Protein Isolates", *Food Hydrocolloids*, Vol. 22, pp. 560-567.
- Wardlaw G M (2004), *Perspectives in Nutrition*, 6th Edition, McGraw Hill Companies, New York, USA.
- Wikipedia (2010), *Cereal* (October 13), Retrieved October 18, from <http://en.wikipedia.org/wiki/cereal>
- Yusufu PA, Egbunu FA, Egwujeh S ID, Opega G L and Adikwu M O (2013), "Evaluation of Complementary Food Prepared from Sorghum, African Yam Bean and Mango Mesocarp Flour Blends", *Pakistan Journal of Nutrition*, Vol. 12, No. 2, pp. 205-208.

